

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/098890 A2

(51) International Patent Classification⁷: **C07H**

(US). **LIUBIN, Mario, N.** [US/US]; 3014 Los Prados # A310, San Mateo, CA 94403 (US).

(21) International Application Number: PCT/US02/17461

(22) International Filing Date: 3 June 2002 (03.06.2002)

(74) Agents: **BRUNELLE, Jan** et al.; Exelixis, Inc., P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/296,076 5 June 2001 (05.06.2001) US
60/328,605 10 October 2001 (10.10.2001) US
60/357,253 15 February 2002 (15.02.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **EX-ELIXIS, INC.** [US/US]; 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRIEDMAN, Lori** [US/US]; One Bayside Village Place, Unit 212, San Francisco, CA 94107 (US). **PLOWMAN, Gregory, D.** [US/US]; 35 Winding Way, San Carlos, CA 94070 (US). **BELVIN, Marcia** [US/US]; 921 Santa Fe Avenue, Albany, CA 94706 (US). **FRANCIS-LANG, Helen** [GB/US]; 1782 Pacific Avenue, Apt. 2, San Francisco, CA 94109 (US). **LI, Danxi** [CN/US]; 90 Behr Avenue #302, San Francisco, CA 94131 (US). **FUNKE, Roel, P.** [NL/US]; 343 California Avenue, South San Francisco, CA 94080

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MARKs AS MODIFIERS OF THE p53 PATHWAY AND METHODS OF USE

(57) Abstract: Human MARK genes are identified as modulators of the p53 pathway, and thus are therapeutic targets for disorders associated with defective p53 function. Methods for identifying modulators of p53, comprising screening for agents that modulate the activity of MARK are provided.

MARKs AS MODIFIERS OF THE p53 PATHWAY AND METHODS OF USE**REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. provisional patent applications 60/296,076
5 filed 6/5/2001, 60/328,605 filed 10/10/2001, and 60/357,253 filed 2/15/2002. The
contents of the prior applications are hereby incorporated in their entirety.

BACKGROUND OF THE INVENTION

The p53 gene is mutated in over 50 different types of human cancers, including
10 familial and spontaneous cancers, and is believed to be the most commonly mutated gene
in human cancer (Zambetti and Levine, FASEB (1993) 7:855-865; Hollstein, *et al.*,
Nucleic Acids Res. (1994) 22:3551-3555). Greater than 90% of mutations in the p53 gene
are missense mutations that alter a single amino acid that inactivates p53 function.
Aberrant forms of human p53 are associated with poor prognosis, more aggressive tumors,
15 metastasis, and short survival rates (Mitsudomi *et al.*, Clin Cancer Res 2000 Oct;
6(10):4055-63; Koshland, Science (1993) 262:1953).

The human p53 protein normally functions as a central integrator of signals including
DNA damage, hypoxia, nucleotide deprivation, and oncogene activation (Prives, Cell
(1998) 95:5-8). In response to these signals, p53 protein levels are greatly increased with
20 the result that the accumulated p53 activates cell cycle arrest or apoptosis depending on
the nature and strength of these signals. Indeed, multiple lines of experimental evidence
have pointed to a key role for p53 as a tumor suppressor (Levine, Cell (1997) 88:323-331).
For example, homozygous p53 "knockout" mice are developmentally normal but exhibit
nearly 100% incidence of neoplasia in the first year of life (Donehower *et al.*, Nature
25 (1992) 356:215-221).

The biochemical mechanisms and pathways through which p53 functions in normal
and cancerous cells are not fully understood, but one clearly important aspect of p53
function is its activity as a gene-specific transcriptional activator. Among the genes with
known p53-response elements are several with well-characterized roles in either regulation
30 of the cell cycle or apoptosis, including GADD45, p21/Waf1/Cip1, cyclin G, Bax, IGF-
BP3, and MDM2 (Levine, Cell (1997) 88:323-331).

Microtubules have a central role in the regulation of cell shape and polarity during
differentiation, chromosome partitioning at mitosis, and intracellular transport.
Microtubules undergo rearrangements involving rapid transitions between stable and

dynamic states during these processes. Microtubule affinity regulating kinases (MARK) are a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption (Drewes, G., et al. (1997) Cell 89: 297-308).

5 Microtubule affinity regulating kinase 1 (MARK1) is a serine/threonine kinase that phosphorylates microtubule-associated protein tau, leading to disruption of microtubules. It shares 90% amino acid homology with the rat version of MARK1, and demonstrates ubiquitous expression with highest levels in testis and brain (Nagase, T. et al. (2000) DNA Res. 7: 143-150).

10 EMK1 (MARK2) is a serine/threonine protein kinase with two isoforms, which differ by the presence or absence of a 162-bp alternative exon (Espinosa, L. and Navarro, E. (1998) Cytogenet. Cell Genet. 81:278-282). Both human isoforms are coexpressed in a number of cell lines and tissues, with the highest expression found in heart, brain, placenta, skeletal muscle, and pancreas, and at lower levels in lung, liver, and kidney (Inglis, J. et al. (1993) Mammalian Genome 4: 401-403). Due to the physical location of
15 this gene, 11q12-q13, EMK1 is a candidate gene for carcinogenic events (Courseaux, A. et al. (1995) Mammalian Genome 6: 311-312), and has been associated with colon and prostate cancer (Moore, T. M., et al. (2000) J Biol Chem 275:4311-22; Navarro, E., et al. (1999) Biochim Biophys Acta 1450: 254-64).

20 Microtubule affinity regulating kinase 3 (MARK3) was originally identified as a marker (KP78) induced by treatment with DNA damaging agents. The loss of MARK3 was associated with carcinogenesis in the pancreas (Parsa, I. (1988) Cell Growth Differ. 9: 197-208). MARK3 may be involved in cell cycle regulation, and alterations in the MARK3 gene may lead to carcinogenesis. MARK 3 is ubiquitously expressed throughout human tissues, with an additional 3.0 Kb transcript present in the heart (Peng, C. et al.
25 (1998) Cell Growth Differ. 9: 197-208).

MAP/microtubule affinity-regulating kinase like 1 (MARKL1) has two isoforms (Nagase, T. et al. (2001) DNA Res. 8: 85-95), is activated by the beta-catenin/Tcf complex in hepatic cell lines, and may be involved in hepatic carcinogenesis (Kato, T. et al. (2001). Neoplasia 3:4-9).

30 The ability to manipulate the genomes of model organisms such as *Drosophila* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, has direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and

mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Mechler BM et al., 1985 EMBO J 4:1551-1557; Gateff E. 1982 Adv. Cancer Res. 37: 33-74; Watson KL., et al., 1994 J Cell Sci. 18: 19-33; Miklos GL, and Rubin GM. 1996 Cell 86:521-529; Wassarman DA, et al., 1995 Curr Opin Gen Dev 5: 44-50; and Booth DR. 1999 Cancer Metastasis Rev. 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as p53, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including sequence information in referenced Genbank identifier numbers and website references, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the p53 pathway in *Drosophila*, and identified their human orthologs, hereinafter referred to as MARK. The invention provides methods for utilizing these p53 modifier genes and polypeptides to identify candidate therapeutic agents that can be used in the treatment of disorders associated with defective p53 function. Preferred MARK-modulating agents specifically bind to MARK polypeptides and restore p53 function. Other preferred MARK-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress MARK gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

MARK-specific modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with a MARK polypeptide or nucleic acid. In one embodiment, candidate p53 modulating agents are tested with an assay system comprising a MARK polypeptide or nucleic acid. Candidate agents that produce a change in the activity of the assay system relative to controls are identified as candidate p53 modulating agents. The assay system may be cell-based or cell-free. MARK-modulating agents

include MARK related proteins (e.g. dominant negative mutants, and biotherapeutics); MARK-specific antibodies; MARK-specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind MARK or compete with MARK binding target. In one specific embodiment, a small molecule modulator is identified
5 using a kinase assay. In specific embodiments, the screening assay system is selected from a binding assay, an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

In another embodiment, candidate p53 pathway modulating agents are further tested using a second assay system that detects changes in the p53 pathway, such as angiogenic,
10 apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the p53 pathway, such as an angiogenic, apoptotic, or cell
15 proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the p53 pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a MARK polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal
20 predetermined to have a pathology associated the p53 pathway.

DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the p53 pathway in *Drosophila* in which p53 was overexpressed in the wing (Ollmann M, et al., Cell 2000 101: 91-101).
25 The KP78a gene was identified as a modifier of the p53 pathway. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, microtubule affinity regulator kinase (MARK) genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective p53 signaling pathway, such as cancer.

In vitro and in vivo methods of assessing MARK function are provided herein.
30 Modulation of the MARK or their respective binding partners is useful for understanding the association of the p53 pathway and its members in normal and disease conditions and for developing diagnostics and therapeutic modalities for p53 related pathologies. MARK-modulating agents that act by inhibiting or enhancing MARK expression, directly

or indirectly, for example, by affecting a MARK function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. MARK modulating agents are useful in diagnosis, therapy and pharmaceutical development.

5 **Nucleic acids and polypeptides of the invention**

Sequences related to MARK nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) number) as GI#s 9845486 (SEQ ID NO:1), 9845488 (SEQ ID NO:2), 18578044 (SEQ ID NO:3), 14250621 (SEQ ID NO:6), 15042610 (SEQ ID NO:7), 8923921 (SEQ ID NO:8),
 10 17445805 (SEQ ID NO:9), 7959214 (SEQ ID NO:11), 14042208 (SEQ ID NO:12),
 3089348 (SEQ ID NO:13), 4505102 (SEQ ID NO:14), 5714635 (SEQ ID NO:15),
 18448970 (SEQ ID NO:18), 13366083 (SEQ ID NO:19), 14017936 (SEQ ID NO:22), and
 16555377 (SEQ ID NO:23) for nucleic acid, and GI#s 9845487 (SEQ ID NO:24),
 8923922 (SEQ ID NO:25), 3089349 (SEQ ID NO:26), 4505103 (SEQ ID NO:27),
 15 13366084 (SEQ ID NO:28) and 13899225 (SEQ ID NO:29) for polypeptides.
 Additionally, nucleic acid sequences of SEQ ID NOs:4, 5, 16, 17, 20, 21, and novel nucleic acid sequence of SEQ ID NO:10 can also be used in the invention.

MARKs are kinase proteins with kinase and UBA/TS-N domains. The term "MARK polypeptide" refers to a full-length MARK protein or a functionally active fragment or
 20 derivative thereof. A "functionally active" MARK fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type MARK protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of MARK proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in
 25 Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of a MARK, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2;
 30 <http://pfam.wustl.edu>). For example, the protein kinase domains of MARKs from GI#s 9845487 (SEQ ID NO:24), 8923922 (SEQ ID NO:25), 4505103 (SEQ ID NO:27), and 13899225 (SEQ ID NO:29) is located at approximately amino acid residues 20 to 271, 60 to 311, 56 to 307, and 59 to 310, respectively (PFAM 00069). Further, the ubiquitin associated (UBA/TS-N) domains of MARKs from GI#s 9845487 (SEQ ID NO:24),

8923922 (SEQ ID NO:25), 4505103 (SEQ ID NO:27) and 13899225 (SEQ ID NO:29) is located at approximately amino acid residues 291 to 330, 331 to 370, 327 to 366, and 330 to 369, respectively (PFAM 00627). Methods for obtaining MARK polypeptides are also further described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of any one of SEQ ID NOs:24, 25, 26, 27, 28, or 29 (a MARK). In further preferred embodiments, the fragment comprises the entire kinase (functionally active) domain.

10 The term "MARK nucleic acid" refers to a DNA or RNA molecule that encodes a MARK polypeptide. Preferably, the MARK polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with MARK.

15 Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST

20 (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences

25 from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in

30 one species, such as *Drosophila*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the

subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410; <http://blast.wustl.edu/blast/README.html>) with all the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute <http://www.ebi.ac.uk/MPsrch/>; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap

extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

- Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of any of SEQ ID NOs:1 through 23.
- 5 The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments,
- 10 a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of any one of SEQ ID NOs:1 through 23 under stringent hybridization conditions that comprise: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X
- 15 Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.2X SSC and 0.1% SDS (sodium dodecyl sulfate).
- 20 In other embodiments, moderately stringent hybridization conditions are used that comprise: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl
- 25 (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

- Alternatively, low stringency conditions can be used that comprise: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM
- 30 sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of MARK Nucleic Acids and Polypeptides

MARK nucleic acids and polypeptides, useful for identifying and testing agents that modulate MARK function and for other applications related to the involvement of MARK in the p53 pathway. MARK nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of a MARK protein for assays used to assess MARK function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) *Protein Expression: A Practical Approach*, Oxford University Press Inc., New York 1999; Stanbury PF et al., *Principles of Fermentation Technology*, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) *Protein Purification Protocols*, Humana Press, New Jersey, 1996; Coligan JE et al, *Current Protocols in Protein Science* (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant MARK is expressed in a cell line known to have defective p53 function (*e.g.* SAOS-2 osteoblasts, H1299 lung cancer cells, C33A and HT3 cervical cancer cells, HT-29 and DLD-1 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding a MARK polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native MARK gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus);

microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. A host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the MARK gene product, the expression vector can comprise a promoter operably linked to a MARK gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the MARK gene product based on the physical or functional properties of the MARK protein in *in vitro* assay systems (*e.g.* immunoassays).

The MARK protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (*i.e.* it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the MARK gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis, cite purification reference). Alternatively, native MARK proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of MARK or other genes associated with the p53 pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter MARK expression may be used in *in vivo* assays to test for activity of a candidate p53 modulating agent, or to further assess the role of MARK in a p53 pathway process such as apoptosis or cell proliferation.

Preferably, the altered MARK expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal MARK expression. The genetically modified animal may additionally have altered p53 expression (e.g. p53 knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice), cows, horses, goats, sheep, pigs, dogs and cats. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, i.e. mosaic animals (see, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) or stably integrated into its germ line DNA (i.e., in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster et al., Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Pat. No. 4,873,191 by Wagner et al., and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous MARK gene that results in a decrease of MARK function, preferably such that MARK expression is

undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse MARK gene is used to construct a homologous recombination vector suitable for altering an endogenous MARK gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) Scan J Immunol 40:257-264; Declerck PJ *et al.*, (1995) J Biol Chem. 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the MARK gene, e.g., by introduction of additional copies of MARK, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the MARK gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment,

both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X et al (2000) Nat Genet 25:83-6).

5 The genetically modified animals can be used in genetic studies to further elucidate the p53 pathway, as animal models of disease and disorders implicating defective p53 function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered MARK function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that
10 receive placebo treatment, and/or animals with unaltered MARK expression that receive candidate therapeutic agent.

In addition to the above-described genetically modified animals having altered MARK function, animal models having defective p53 function (and otherwise normal MARK function), can be used in the methods of the present invention. For example, a p53
15 knockout mouse can be used to assess, *in vivo*, the activity of a candidate p53 modulating agent identified in one of the *in vitro* assays described below. p53 knockout mice are described in the literature (Jacks et al., Nature 2001;410:1111-1116, 1043-1044; Donehower *et al.*, supra). Preferably, the candidate p53 modulating agent when administered to a model system with cells defective in p53 function, produces a detectable
20 phenotypic change in the model system indicating that the p53 function is restored, i.e., the cells exhibit normal cell cycle progression.

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate
25 the function of MARK and/or the p53 pathway. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the p53 pathway, as well as in further analysis of the MARK protein and its contribution to the p53 pathway. Accordingly, the invention also provides methods for modulating the p53 pathway comprising the step of specifically modulating MARK activity by administering a MARK-
30 interacting or -modulating agent.

In a preferred embodiment, MARK-modulating agents inhibit or enhance MARK activity or otherwise affect normal MARK function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a further preferred embodiment, the candidate p53 pathway- modulating agent specifically

modulates the function of the MARK. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the MARK polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the MARK. The term also encompasses modulating agents that alter the interaction of the MARK with a binding partner or substrate (e.g. by binding to a binding partner of a MARK, or to a protein/binding partner complex, and inhibiting function).

Preferred MARK-modulating agents include small molecule compounds; MARK-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

Small molecule modulators

Small molecules, are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the MARK protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for MARK-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the p53 pathway. The activity of candidate small molecule

modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific MARK-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the p53 pathway and related disorders, as well as in validation assays for other MARK-modulating agents. In a preferred embodiment, MARK-interacting proteins affect normal MARK function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, MARK-interacting proteins are useful in detecting and providing information about the function of MARK proteins, as is relevant to p53 related disorders, such as cancer (e.g., for diagnostic means).

An MARK-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with a MARK, such as a member of the MARK pathway that modulates MARK expression, localization, and/or activity. MARK-modulators include dominant negative forms of MARK-interacting proteins and of MARK proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous MARK-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

An MARK-interacting protein may be an exogenous protein, such as a MARK-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). MARK antibodies are further discussed below.

In preferred embodiments, a MARK-interacting protein specifically binds a MARK protein. In alternative preferred embodiments, a MARK-modulating agent binds a MARK substrate, binding partner, or cofactor.

5 *Antibodies*

In another embodiment, the protein modulator is a MARK specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify MARK modulators. The antibodies can also be used in dissecting the portions of the MARK pathway responsible for various cellular responses and in the general processing and maturation of the MARK.

10 Antibodies that specifically bind MARK polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of MARK polypeptide, and more preferably, to human MARK. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab
15 fragments, F(ab')₂ fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of MARK which are particularly antigenic can be selected, for example, by routine screening of MARK polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Natl.
20 Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89; Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence shown in any of SEQ ID NOs:24, 25, 26, 27, 28, or 29. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles
25 and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of MARK or substantially purified fragments thereof. If MARK fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of a MARK protein. In a particular embodiment, MARK-specific antigens and/or immunogens are
30 coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of MARK-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding MARK polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

- 5 Chimeric antibodies specific to MARK polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., 10 Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies 15 into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 20 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

- MARK-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 25 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

- Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As 30 used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that

express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Nucleic Acid Modulators

Other preferred MARK-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit MARK activity. Preferred nucleic acid modulators interfere with the function of the MARK nucleic acid such as DNA replication, transcription, translocation of the MARK RNA to the site of protein translation, translation of protein from the MARK RNA, splicing of the MARK RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the MARK RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to a MARK mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. MARK-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred MARK nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al*., Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, a MARK-specific nucleic acid modulator is used in an assay to further elucidate the role of the MARK in the p53 pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, a MARK-specific antisense oligomer is used as a therapeutic agent for treatment of p53-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of MARK activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the MARK nucleic acid or protein. In general, secondary assays further assess the activity of a MARK modulating agent identified by a primary assay and may confirm that the modulating agent affects MARK in a manner relevant to the p53 pathway. In some cases, MARK modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising a MARK polypeptide with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates MARK activity, and hence the p53 pathway.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

5 For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead
10 cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding),
15 transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular
20 molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of MARK and any auxiliary proteins demanded by the particular assay. Appropriate
1 methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity
25 and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when MARK-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the MARK protein may be assayed by various known methods
30 such as substrate processing (*e.g.* ability of the candidate MARK-specific binding agents to function as negative effectors in MARK-expressing cells), binding equilibrium constants (usually at least about 10^7 M⁻¹, preferably at least about 10^8 M⁻¹, more preferably at least about 10^9 M⁻¹), and immunogenicity (*e.g.* ability to elicit MARK specific antibody

in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a MARK polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The MARK polypeptide can be full length or a fragment thereof that retains functional MARK activity. The MARK polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The MARK polypeptide is preferably human MARK, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of MARK interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has MARK-specific binding activity, and can be used to assess normal MARK gene function.

Suitable assay formats that may be adapted to screen for MARK modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (e.g., Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate MARK and p53 pathway modulators (e.g. U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); U.S. Pat. No. 6,020,135 (p53 modulation), among others). Specific preferred assays are described in more detail below.

Kinase assays. In some preferred embodiments the screening assay detects the ability of the test agent to modulate the kinase activity of a MARK polypeptide. In further embodiments, a cell-free kinase assay system is used to identify a candidate p53 modulating agent, and a secondary, cell-based assay, such as an apoptosis or hypoxic induction assay (described below), may be used to further characterize the candidate p53 modulating agent. Many different assays for kinases have been reported in the literature

and are well known to those skilled in the art (e.g. U.S. Pat. No. 6,165,992; Zhu et al., Nature Genetics (2000) 26:283-289; and WO0073469). Radioassays, which monitor the transfer of a gamma phosphate are frequently used. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from gamma -³³P ATP to a biotinylated peptide substrate; the substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand.

Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133).

Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). An apoptosis assay system may comprise a cell that expresses a MARK, and that optionally has defective p53 function (e.g. p53 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate p53 modulating agents. In some embodiments of the invention, an apoptosis

assay may be used as a secondary assay to test a candidate p53 modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether MARK function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express MARK relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the MARK plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen, J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed with MARK are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW *et al.* (1986) *Int J Radiat Biol Relat Stud Phys Chem Med* 49:237-55). Cells transfected with a MARK may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson).

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses a MARK, and that optionally has defective p53 function (e.g. p53 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate p53 modulating agents. In some embodiments of

the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate p53 modulating agents that is initially identified using another assay system such as a cell-free kinase assay system. A cell proliferation assay may also be used to test whether MARK function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express MARK relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the MARK plays a direct role in cell proliferation or cell cycle.

10 **Angiogenesis.** Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses a MARK, and that optionally has defective p53 function (e.g. p53 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate p53 modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate p53 modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether MARK function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express MARK relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the MARK plays a direct role in angiogenesis.

30 **Hypoxic induction.** The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells

transfected with MARK in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses a MARK, and that optionally has a mutated p53 (e.g. p53 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate p53 modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate p53 modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether MARK function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express MARK relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the MARK plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents.

Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., *Bioconjug Chem.* 2001 May-Jun;12(3):346-53).

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the MARK protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting MARK-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance MARK gene expression, preferably mRNA expression. In general, expression analysis comprises comparing MARK expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express MARK) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that MARK mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, *Biotechniques* (1999) 26:112-125; Kallioniemi OP, *Ann Med* 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A *Curr Opin Biotechnol* 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the MARK protein or

specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

Secondary Assays

5 Secondary assays may be used to further assess the activity of MARK-modulating agent identified by any of the above methods to confirm that the modulating agent affects MARK in a manner relevant to the p53 pathway. As used herein, MARK-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a
10 modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with MARK.

 Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express MARK) in the presence and absence of the candidate modulator. In general, such assays test whether
15 treatment of cells or animals with a candidate MARK-modulating agent results in changes in the p53 pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use "sensitized genetic backgrounds", which, as used herein, describe cells or animals engineered for altered expression of genes in the p53 or interacting pathways.

20

Cell-based assays

 Cell based assays may use a variety of mammalian cell lines known to have defective p53 function (*e.g.* SAOS-2 osteoblasts, H1299 lung cancer cells, C33A and HT3 cervical cancer cells, HT-29 and DLD-1 colon cancer cells, among others, available from
25 American Type Culture Collection (ATCC), Manassas, VA). Cell based assays may detect endogenous p53 pathway activity or may rely on recombinant expression of p53 pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

30

Animal Assays

 A variety of non-human animal models of normal or defective p53 pathway may be used to test candidate MARK modulators. Models for defective p53 pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express

or lack expression in) genes involved in the p53 pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, p53 pathway activity is assessed by monitoring
5 neovascularization and angiogenesis. Animal models with defective and normal p53 are used to test the candidate modulator's affect on MARK in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents,
10 such as bFGF and VEGF, or with human tumor cells which over-express the MARK. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel®
15 pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on MARK is assessed via tumorigenicity assays. In one example, xenograft human tumors are implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either
20 from a pre-existing tumor or from *in vitro* culture. The tumors which express the MARK endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 μ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus
25 administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors maybe utilized for biomarker identification or further analyses. For
30 immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

Diagnostic and therapeutic uses

Specific MARK-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the p53 pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the p53 pathway in a cell, preferably a cell pre-determined to have defective p53 function, comprising the step of administering an agent to the cell that specifically modulates MARK activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the p53 function is restored, i.e., for example, the cell undergoes normal proliferation or progression through the cell cycle.

The discovery that MARK is implicated in p53 pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the p53 pathway and for the identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether MARK expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective p53 signaling that express a MARK, are identified as amenable to treatment with a MARK modulating agent. In a preferred application, the p53 defective tissue overexpresses a MARK relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial MARK cDNA sequences as probes, can determine whether particular tumors express or overexpress MARK. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of MARK expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the MARK oligonucleotides, and antibodies directed against a MARK, as described above for: (1) the detection of the presence of MARK gene mutations, or the detection of either over- or under-expression of MARK mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of MARK gene

product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by MARK.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease in a patient, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for MARK expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of disease. Preferably, the disease is cancer, most preferably a cancer as shown in TABLE 1. The probe may be either DNA or protein, including an antibody.

10 EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

I. Drosophila p53 screen

15 The *Drosophila* p53 gene was overexpressed specifically in the wing using the vestigial margin quadrant enhancer. Increasing quantities of *Drosophila* p53 (titrated using different strength transgenic inserts in 1 or 2 copies) caused deterioration of normal wing morphology from mild to strong, with phenotypes including disruption of pattern and polarity of wing hairs, shortening and thickening of wing veins, progressive crumpling of the wing and appearance of dark “death” inclusions in wing blade. In a screen designed to identify enhancers and suppressors of *Drosophila* p53, homozygous females carrying two copies of p53 were crossed to 5663 males carrying random insertions of a piggyBac transposon (Fraser M *et al.*, Virology (1985) 145:356-361). Progeny containing insertions were compared to non-insertion-bearing sibling progeny for enhancement or suppression of the p53 phenotypes. Sequence information surrounding the piggyBac insertion site was used to identify the modifier genes. Modifiers of the wing phenotype were identified as members of the p53 pathway. kp78a was a suppressor of the wing phenotype. Human orthologs of the modifiers are referred to herein as MARK.

BLAST analysis (Altschul et al., *supra*) was employed to identify Targets from *Drosophila* modifiers. For example, representative sequences from MARK GI# 9845487 (SEQ ID NO:24), GI# 8923922(SEQ ID NO:25), GI# 4505103 (SEQ ID NO:27), and GI#13899225 (SEQ ID NO:29) share 43%, 65%, 65% and 45% amino acid identity, respectively, with the *Drosophila* kp78a.

Various domains, signals, and functional subunits in proteins were analyzed using the PSORT (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6; Kenta Nakai, Protein sorting signals and prediction of subcellular localization, Adv. Protein Chem. 54, 277-344 (2000)), PFAM (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2; <http://pfam.wustl.edu>), SMART (Ponting CP, et al., SMART: identification and annotation of domains from signaling and extracellular protein sequences. Nucleic Acids Res. 1999 Jan 1;27(1):229-32), TM-HMM (Erik L.L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998), and clust (Remm M, and Sonnhammer E. Classification of transmembrane protein families in the *Caenorhabditis elegans* genome and identification of human orthologs. Genome Res. 2000 Nov;10(11):1679-89) programs. For example, the protein kinase domains of MARKs from GI#s 9845487 (SEQ ID NO:24), 8923922 (SEQ ID NO:25), 4505103 (SEQ ID NO:27), and 13899225 (SEQ ID NO:29) is located at approximately amino acid residues 20 to 271, 60 to 311, 56 to 307, and 59 to 310, respectively (PFAM 00069). Further, the ubiquitin associated (UBA/TS-N) domains of MARKs from GI#s 9845487 (SEQ ID NO:24), 8923922 (SEQ ID NO:25), 4505103 (SEQ ID NO:27), and 13899225 (SEQ ID NO:29) is located at approximately amino acid residues 291 to 330, 331 to 370, 327 to 366, and 330 to 369, respectively (PFAM 00627). Still further, the kinase associated domains from MARKs of GI#s 9845487 (SEQ ID NO:24), 8923922 (SEQ ID NO:25), and 4505103 (SEQ ID NO:27) are located at approximately amino acid residues 696 to 745, 746 to 795, and 664 to 713, respectively (PFAM 02149).

25

II. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled MARK peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of MARK activity.

30

III. High-Throughput In Vitro Binding Assay.

³³P-labeled MARK peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-
5 avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate p53 modulating agents.

10 IV. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the MARK proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once
15 with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is
20 incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated
25 antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

V. Kinase assay

A purified or partially purified MARK is diluted in a suitable reaction buffer, e.g., 50
30 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 μ g/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μ l. The

reaction is initiated by the addition of ^{33}P -gamma-ATP (0.5 $\mu\text{Ci/ml}$) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation (Mg^{2+} or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VI. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, and Ambion.

TaqMan analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/ μl . Single stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA, <http://www.appliedbiosystems.com/>).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan protocols, and the following criteria: a) primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product.

Taqman reactions were carried out following manufacturer's protocols, in 25 μl total volume for 96-well plates and 10 μl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis was prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in

appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched normal tissues from the same patient. A gene was considered overexpressed in a tumor when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample.

Results are shown in Table 1. Data presented in bold indicate that greater than 50% of tested tumor samples of the tissue type indicated in row 1 exhibited over expression of the gene listed in column 1, relative to normal samples. Underlined data indicates that between 25% to 49% of tested tumor samples exhibited over expression. A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

Table 1

	<u>breast</u>		<u>colon</u>		<u>lung</u>		<u>ovary</u>	
GI#9845486 (SEQ ID NO:1)	7	11	<u>8</u>	<u>30</u>	8	13	5	7
GI#9845488 (SEQ ID NO:2)	1	11	4	30	0	13	1	7
GI#8923921 (SEQ ID NO:8)	2	11	7	30	<u>6</u>	<u>13</u>	0	7
GI#3089348 (SEQ ID NO:13)	2	11	2	30	0	13	<u>2</u>	<u>7</u>
GI#13366083 (SEQ ID NO:19)	2	11	2	30	<u>5</u>	<u>13</u>	1	7

WHAT IS CLAIMED IS:

1. A method of identifying a candidate p53 pathway modulating agent, said method comprising the steps of:
 - 5 (a) providing an assay system comprising a purified MARK polypeptide or nucleic acid or a functionally active fragment or derivative thereof;
 - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
 - (c) detecting a test agent-biased activity of the assay system, wherein a difference
 - 10 between the test agent-biased activity and the reference activity identifies the test agent as a candidate p53 pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the MARK polypeptide.
- 15 3. The method of Claim 2 wherein the cultured cells additionally have defective p53 function.
4. The method of Claim 1 wherein the assay system includes a screening assay
- 20 comprising a MARK polypeptide, and the candidate test agent is a small molecule modulator.
5. The method of Claim 4 wherein the assay is a kinase assay.
- 25 6. The method of Claim 1 wherein the assay system is selected from the group consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay
- 30 comprising a MARK polypeptide and the candidate test agent is an antibody.
8. The method of Claim 1 wherein the assay system includes an expression assay comprising a MARK nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
- 5 11. The method of Claim 1 additionally comprising:
(d) administering the candidate p53 pathway modulating agent identified in (c) to a model system comprising cells defective in p53 function and, detecting a phenotypic change in the model system that indicates that the p53 function is restored.
- 10 12. The method of Claim 11 wherein the model system is a mouse model with defective p53 function.
13. A method for modulating a p53 pathway of a cell comprising contacting a cell
15 defective in p53 function with a candidate modulator that specifically binds to a MARK polypeptide comprising an amino acid sequence selected from group consisting of SEQ ID NOs:24, 25, 26, 27, 28, and 29, whereby p53 function is restored.
14. The method of claim 13 wherein the candidate modulator is administered to a
20 vertebrate animal predetermined to have a disease or disorder resulting from a defect in p53 function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
- 25 16. The method of Claim 1, comprising the additional steps of:
(d) providing a secondary assay system comprising cultured cells or a non-human animal expressing MARK ,
(e) contacting the secondary assay system with the test agent of (b) or an agent
30 derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and
(f) detecting an agent-biased activity of the second assay system,

wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate p53 pathway modulating agent,
and wherein the second assay detects an agent-biased change in the p53 pathway.

5

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

10

18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a p53 pathway gene.

15

20. A method of modulating p53 pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a MARK polypeptide or nucleic acid.

21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the p53 pathway.

20

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

23. A method for diagnosing a disease in a patient comprising:
25 (a) obtaining a biological sample from the patient;
(b) contacting the sample with a probe for MARK expression;
(c) comparing results from step (b) with a control;
(d) determining whether step (c) indicates a likelihood of disease.

30

24. The method of claim 23 wherein said disease is cancer.

25. The method according to claim 24, wherein said cancer is a cancer as shown in Table 1 as having >25% expression level.

SEQUENCE LISTING

<110> EXELIXIS, INC.

<120> MARKs AS MODIFIERS OF THE p53 PATHWAY AND METHODS OF USE

<130> EX02-088C-PC

<150> US 60/296,076

<151> 2001-06-05

<150> US 60/328,605

<151> 2001-10-10

<150> US 60/357,253

<151> 2002-02-15

<160> 29

<170> PatentIn version 3.1

<210> 1

<211> 2946

<212> DNA

<213> Homo sapiens

<400> 1

tcctggaatt gcacgcgctt cctgaccacc aggctctggc ccttgagaag ccagcggggc	60
tttgtccctg ttgctctcct tgccaaaccc agtctctctg ctagtgggtg tttcggttgc	120
gacaccgtcc aggttcccag gcaggaaccg ctggccttg ctgcttagct acttttcact	180
gaggaggtgg tggaaggtgt cgcctgctct ggctgagtaa ggggtggctgg ctgagccggc	240
agcccccgcc ctaggcctgg ctcttcccgg cctctgtact ttgccctcgc tgcctgacag	300
gttctgctgt gggctctgct gaatggaagt cgctggtagt cttttccct ttctccagtc	360
ggccacacctt gggacacctt gactccaagc ccagcagtaa gtccaacatg attcggggcc	420
gcaactcagc cacctctgct gatgagcagc cccacattgg aaactaccgg ctctcaaga	480
ccattggcaa gggtaatttt gccaaaggtga agttggcccg acacatcctg actgggaaag	540
aggtagctgt gaagatcatt gacaagactc aactgaactc ctccagcctc cagaaactat	600
tccgcgaagt aagaataatg aaggttttga atcatccaa catagttaaa ttatttgaag	660
tgattgagac tgagaaaacg ctctaccttg tcatggagta cgctagtggc ggagaggtat	720
ttgattacct agtggctcat ggcaggatga aagaaaaaga ggctcgagcc aaattccgcc	780
agatagtgtc tgctgtgcag tactgtcacc agaagtttat tgtccataga gacttaaagg	840
cagaaaacct gctcttggat gctgatatga acatcaagat tgcagacttt ggcttcagca	900
atgaattcac ctttgggaac aagctggaca ccttctgtgg cagtccccct tatgctgccc	960
cagaactctt ccagggcaaa aaatatgatg gacccgaggt ggatgtgtgg agcctaggag	1020
ttatcctcta tacactggtc agcggatccc tgcccttttga tggacagaac ctcaaggagc	1080

tgccgggaacg ggtactgagg gggaaatacc gtattccatt ctacatgtcc acggactgtg	1140
aaaacctgct taagaaatit ctcatcctta atcccagcaa gagaggcact ttagagcaaa	1200
tcatgaaaga tcgatggatg aatgtgggtc acgaagatga tgaactaaag ccttacgtgg	1260
agccactccc tgactacaag gacccccggc ggacagagct gatgggtgtcc atggggtata	1320
cacgggaaga gatccaggac tcgctgggtg gccagagata caacgagggt atgggccacct	1380
atctgctcct gggctacaag agctccgagc tgggaaggcg caccatcacc ctgaaacccc	1440
ggccttcagc tgatctaacc aatagcagcg cccaattccc atcccacaag gtacagcgaa	1500
gcgtgtcggc caatcccaag cagcggcgct tcagcgacca ggctgggtcct gccattccca	1560
cctctaattc ttactctaag aagactcaga gtaacaacgc agaaaataag cggcctgagg	1620
aggaccggga gtcagggcgg aaagccagca gcacagccaa ggtgcctgcc agccccctgc	1680
ccggtctgga gaggaagaag accaccccaa cccctccac gaacagcgtc ctctccacca	1740
gcacaaatcg aagcaggaat tccccacttt tggagcgggc cagcctcggc caggcctcca	1800
tccagaatgg caaagacagc ctaaccatgc caggggtccg ggctccacg gcttctgctt	1860
ctgccgcagt ctctcgggc cggccccgcc agcaccagaa atccatgtcg gctccgtgc	1920
acccaacaa ggctcttggg ctgccccca cggagagtaa ctgtgagggt cgcgggcca	1980
gcacagcccc ccagcgtgtc cctgttgct ccccatccgc ccacaacatc agcagcagt	2040
gtggagcccc agaccgaact aacttcccc ggggtgtgtc cagccgaagc accttccatg	2100
ctgggcagct ccgacagggt cgggaccagc agaatttgcc ctacgggtgtg accccagcct	2160
ctccctctgg ccacagccag ggccggcggg gggcctctgg gagcatcttc agcaagtcca	2220
cctccaagtt tgtacgcagg aacctgaatg aacctgaaag caaagaccga gtggagacgc	2280
tcagacctca cgtggtgggc agtggcggca acgacaaaga aaaggaagaa ttctgggagg	2340
ccaagccccg ctccctccgc ttcacgtgga gtatgaagac cacgagctcc atggagccca	2400
acgagatgat gcgggagatc cgcaagggtg tggacgcgaa cagctgccag agcagcgtgc	2460
atgagaagta catgtgtgtg tgcattgcac gcacgcggg ccacgaggac ttcgtgcagt	2520
gggagatgga ggtgtgcaaa ctgcccgggc tctctctcaa cggggttcga ttttaagcga	2580
tatcgggcac ctccatggc ttcaaaaaca ttgcctccaa aatagccaac gagctgaagc	2640
tttaacaggc tgccaggagc gggggcggcg ggggcgggccc agctggacgg gctgccggcc	2700
gtgcgccgcc ccacctgggc gagactgcag cgatggattg gtgtgtctcc ctgctggcac	2760
ttctcccctc cctggccctt ctcaagttttc tcccacattc acccctgccc agagattccc	2820
ccttctcttc tcccctactg gaggcaaagg aaggggaggg tggatggggg ggcagggctc	2880
cccctcggtc ctgcggttgc acagagtatt tcgcctaaac caagaaattt ttattacca	2940

aaaaga

2946

<210> 2

<211> 2784

<212> DNA

<213> Homo sapiens

<400> 2

tcctggaatt gcacgcgctt cctgaccacc aggcctctggc ccttgagaag ccagcggggc	60
tttgtccctg ttgctctcct tgccaaaccc agtctctctg ctagtgggtgg tttcgggtgc	120
gacaccgtcc aggttcccag gcaggaaccg ctccggcctgg ctgcttagct acttttcact	180
gaggaggtgg tggaagggtg cgctgctct ggctgagtaa gggtaggtgg ctgagccggc	240
agcccccgcc ctaggcctgg ctcttcccgg cctctgtact ttgccctcgc tgcctgacag	300
gttctgctgt gggctctgct gaatggaagt cgctggtagt ccttttccct ttctccagtc	360
ggcccacctt gggacacctt gactccaagc ccagcagtaa gtccaacatg attcggggcc	420
gcaactcagc cacctctgct gatgagcagc cccacattgg aaactaccgg ctccctcaaga	480
ccattggcaa gggtaatttt gccaaaggtga agttggcccg acacatcctg actgggaaag	540
aggtagctgt gaagatcatt gacaagactc aactgaactc ctccagcctc cagaaactat	600
tccgcgaagt aagaataatg aaggttttga atcatcccaa catagttaaa ttatttgaag	660
tgattgagac tgagaaaacg ctctaccttg tcatggagta cgctagtggc ggagaggtat	720
ttgattacct agtggctcat ggcaggatga aagaaaaaga ggctcgagcc aaattccgcc	780
agatagtgtc tgctgtgcag tactgtcacc agaagtttat tgtccataga gacttaaagg	840
cagaaaacct gctcttggat gctgatatga acatcaagat tgcagacttt ggcttcagca	900
atgaattcac ctttgggaac aagctggaca ccttctgtgg cagtccccct tatgctgccc	960
cagaactctt ccagggcaaa aaatatgatg gacccgaggt ggatgtgtgg agcctaggag	1020
ttatcctcta tacactggtc agcggatccc tgccttttga tggacagaac ctcaaggagc	1080
tgccgggaacg ggtactgagg gggaaatacc gtattccatt ctacatgtcc acggactgtg	1140
aaaacctgct taagaaattt ctcatcctta atcccagcaa gagaggcact ttagagcaaa	1200
tcatgaaaga tcgatggatg aatgtgggtc acgaagatga tgaactaaag ccttacgtgg	1260
agccactccc tgactacaag gacccccggc ggacagagct gatgggtgtcc atgggttata	1320
cacgggaaga gatccaggac tcgctgggtg gccagagata caacgaggtg atggccacct	1380
atctgtctct gggctacaag agctccgagc tggaaaggcga caccatcacg ctgaaacccc	1440
ggccttcagc tgatctaacc aatagcagcg cccaattccc atcccacaag gtacagcgaa	1500
gcgtgtcggc caatcccaag cagcggcgct tcagcgacca ggctggctct gccattccca	1560

```

cctctaattc ttactctaag aagactcaga gtaacaacgc agaaaataag cggcctgagg 1620
aggaccggga gtcagggcgg aaagccagca gcacagccaa ggtgcctgcc agccccctgc 1680
ccggtctgga gaggaagaag accaccccaa cccctccac gaacagcgtc ctctccacca 1740
gcacaaatcg aagcaggaat tccccacttt tggagcgggc cagcctcggc caggcctcca 1800
tccagaatgg caaagacagc acagccccc agcgtgtccc tgttgctcc ccatccgcc 1860
acaacatcag cagcagtggg ggagccccag accgaactaa ctcccccg gggtgtgtcca 1920
gccgaagcac ctccatgct gggcagctcc gacaggtgcg ggaccagcag aatttgccct 1980
acggtgtgac ccagcctct cctctggcc acagccaggg ccggcggggg gcctctggga 2040
gcatcttcag caagttcacc tccaagtttg tacgcaggaa cctgaatgaa cctgaaagca 2100
aagaccgagt ggagacgctc agacctcacg tgggtggcag tggcggcaac gacaaagaaa 2160
aggaagaatt tcgggagggc aagccccgct ccctccgctt cagctggagt atgaagacca 2220
cgagctccat ggagcccaac gagatgatgc gggagatccg caaggtgctg gacgcgaaca 2280
gctgccagag cgagctgcat gagaagtaca tgctgctgtg catgcacggc acgccgggcc 2340
acgaggactt cgtgcagtgg gagatggagg tgtgcaaact gccgcggctc tctctcaacg 2400
gggttcgatt taagcggata tcgggcacct ccatggcctt caaaaacatt gcctccaaaa 2460
tagccaacga gctgaagctt taacaggctg ccaggagcgg gggcggcggg ggcgggccag 2520
ctggacgggc tgccggccgt gcgccgcccc acctgggcga gactgcagcg atggattggt 2580
gtgtctccct gctggcactt ctccccctcc tggcccttct cagttttctc ccacattcac 2640
ccctgccag agattcccc ttctctctc ccctactgga ggcaaaggaa ggggaggggtg 2700
gatggggggg cagggctccc cctcgtact gcggttgac agagtatttc gcctaaacca 2760
agaaatTTTT tattaccaa aaga 2784

```

```

<210> 3
<211> 3103
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (2941)..(2941)
<223> "n" is A, C, G, or T

```

```

<400> 3
cgggtgggtggc ggccatgttg ggagcagcag gtccggcggc ggctgcctgt gtgccggggc 60
cggagcagtg ccgctgaggg caggggagga gcgaggcagg cggccggctg cggcggcaga 120
gagtaggcgg agcggcgcgg cccggccgaa aggcggcaca gccagccgg gggtcggggg 180
ggtgcggtcc ggagccgctc ggagccggcg cggcctagcc cgagcggcgc atccccgggc 240

```

tggcgtgagc ggctgcccgg cctccccgca cccccggccg gggcccatgc ggcgggtgct	300
cctgctgtga gaagccccgc cgggccgggc tccgcgcctt cccttccctc ccttccctcca	360
agctttctcg ttccttcccc cgagataccg ggcgccatgtc cagcgctcgg acccccctac	420
ccacgctgaa cgagaggggac acggagcagc ccaccttggg acaccttgac tccaagccca	480
gcagtaagtc caacatgatt cggggccgca actcagccac ctctgctgat gagcagcccc	540
acattggaaa ctaccggctc ctcaagacca ttggcaaggg taatttttggc aaggtgaagt	600
tggcccgaca catcctgact gggaaagagg tagctgtgaa gatcattgac aagactcaac	660
tgaactcctc cagcctccag aaactattcc gcgaagtaag aataatgaag gttttgaatc	720
atcccaacat agttaaatta tttgaagtga ttgagactga gaaaacgctc taccttgtca	780
tggagtacgc tagtggcgga gaggtatttg attacctagt ggctcatggc aggatgaaag	840
aaaaagaggc tcgagccaaa ttccgccaga tagtgtctgc tgtgcagtac tgtcaccaga	900
agtttattgt ccatagagac ttaaaggcag aaaacctgct cttggatgct gatatgaaca	960
tcaagattgc agactttggc ttcagcaatg aattcacctt tgggaacaag ctggacacct	1020
tctgtggcag tcccccttat gctgccccag aactcttcca gggcaaaaaa tatgatggac	1080
ccgaggtgga tgtgtggagc ctaggagtta tctctatac actgggtcagc ggatccctgc	1140
cttttgatgg acagaacctc aaggagctgc gggaaacgggt actgaggggga aaataccgta	1200
ttccattcta catgtccacg gactgtgaaa acctgcttaa gaaatttctc attcttaatc	1260
ccagcaagag aggcacttta gagcaaatca tgaaagatcg atggatgaat gtgggtcacg	1320
aagatgatga actaaagcct tacgtggagc cactccctga ctacaaggac ccccgccgga	1380
cagagctgat ggtgtccatg ggttatacac gggaagagat ccaggactcg ctgggtgggccc	1440
agagatacaa cgaggtgatg gccacctatc tgctcctggg ctacaagagc tccgagctgg	1500
aaggcgacac catcaccttg aaaccccggc cttcagctga tctgaccaat agcagcgccc	1560
catccccatc ccacaaggta cagcgcagcg tgtcggccaa tcccaagcag cggcgcttca	1620
gcgaccaggc tggctctgcc attcccacct ctaattctta ctctaagaag actcagagta	1680
acaacgcaga aaataagcgg cctgaggagg accgggagtc agggcggaaa gccagcagca	1740
cagccaaggt gcctgccagc cccctgcccg gtctggagag gaagaagacc accccaaccc	1800
cctccacgga acagcgtcct ctccaccagc acaaatcgaa gcaggaatto cccacttttg	1860
gagcggggcca gcctcggcca ggccctccatc cagaatggca aagacagcct aaccatgcca	1920
gggtcccggg cctccacggc ttctgcttct gccgcagtct ctgcggcccg gccccgccag	1980
caccagaaat ccatgtcggc ctccgtgcac cccaacaagg cctctggggt gccccccacg	2040
gagagtaact gtgaggtgcc ggggcccagc acagcccccc agcgtgtccc tgttgccctc	2100

ccatccgccc acaacatcag cagcagtggg ggagccccag accgaaactaa cttcccccg 2160
 ggtgtgtcca gccgaagcac cttccatgct gggcagctcc gacaggtgcg ggaccagcag 2220
 aatttgccct acgggtgtgac ccagcctct ccctctggcc acagccaggg ccggcggggg 2280
 gcctctggga gcattctcag caagttcacc tccaagtgtg tacgcagaaa tctgtctttc 2340
 aggtttgcca gaaggaacct gaatgaacct gaaagcaaag accgagtgga gacgctcaga 2400
 cctcacgtgg tgggcagtgg cggcaacgac aaagaaaagg aagaatttcg ggaggccaag 2460
 ccccgctccc tccgcttcac gtggagtatg aagaccacga gctccatgga gcccaacgag 2520
 atgatcgggg agatccgcaa ggtgctggac gcgaacagct gccagagcga gctgcatgag 2580
 aagtacatgc tgctgtgcat gcacggcacg ccggggccacg aggacttcgt gcagtgggag 2640
 atggaggtgt gcaaactgcc gcggctctct ctcaacgggg ttcgatttaa gcggatatcg 2700
 ggcacctcca tggccttcaa aaacattgcc tccaaaatag ccaacgagct gaagctttaa 2760
 caggctgcca ggagcggggg cggcgggggg cgggccagct ggacgggctg ccggccgctg 2820
 cgccgcccc cctgggcgag actgcagcga tggattggtg tgtctccct gctggcactt 2880
 ctccctccc tggcccttct cagttttctc ttacatgttt gtggggggtg ggagattgtt 2940
 ntccagcacc ccacattcac ccctgccag agattcccc ttctcctctc ccctactgga 3000
 ggcaaaggaa ggggaggggtg gatggggggg cagggctccc cctcggtact gcggttgac 3060
 agagtatttc gcctaaacca agaaattttt tattaccaa aag 3103

<210> 4
 <211> 2086
 <212> DNA
 <213> Homo sapiens

<400> 4
 agtccaacat gattcggggc cgcaactcag ccacctctgc tgatgagcag cccacattg 60
 gaaactaccg gctcctcaag accattggca agggtaattt tgccaagggtg aagttggccc 120
 gacacatcct gactgggaaa gaggtagctg tgaagatcat tgacaagact caactgaact 180
 cctccagcct ccagaaacta ttccgcgaag taagaataat gaagggttttg aatcatccca 240
 acatagttaa attatttgaa gtgattgaga ctgagaaaac gctctacctt gtcattggagt 300
 acgctagtgg cggagaggta tttgattacc tagtggctca tggcaggatg aaagaaaaag 360
 aggctcgagc caaattccgc cagatagtgt ctgctgtgca gtactgtcac cagaagttaa 420
 ttgtccatag agacttaaag gcagaaaacc tgctcttgga tgctgatatg aacatcaaga 480
 ttgcagactt tggcttcagc aatgaattca cctttgggaa caagctggac accttctgtg 540
 gcagtcccc ttatgctgcc ccagaactct tccagggcaa aaaatatgat ggaccggag 600

tggatgtgtg gagcctagga gttatcctct atacactggg cagcggatcc ctgccttttg 660
 atggacagaa cctcaaggag ctgcgggaac ggggtactgag gggaaaatac cgtattccat 720
 tctacatgtc cacggactgt gaaaacctgc ttaagaaatt tctcattctt aatcccagca 780
 agagaggcac tttagagcaa atcatgaaag atcgatggat gaatgtgggt cacgaagatg 840
 atgaactaaa gccttacgtg gagccactcc ctgactacaa ggacccccgg cggacagagc 900
 tgatgggtgtc catgggttat acacgggaag agatccagga ctcgctgggt ggccagagat 960
 acaacgaggt gatggccacc tatctgctcc tgggctacaa gagctccgag ctggaaggcg 1020
 acaccatcac cctgaaaccc cggccttcag ctgatctgac caatagcagc gccccatccc 1080
 catcccacaa ggtacagcgc agcgtgtcgg ccaatcccaa gcagcggcgc ttcagcgcacc 1140
 aggctgggtcc tgccattccc acctctaatt ctactctaa gaagactcag agtaacaacg 1200
 cagaaaataa gcggcctgag gaggaccggg agtcagggcg gaaagccagc agcacagcca 1260
 aggtgcctgc cagccccctg cccggtctgg agaggaagaa gaccacccca accccctcca 1320
 cgaacagcgt cctctccacc agcacaaatc gaagcaggaa tccccactt ttggagcggg 1380
 ccagcctcgg ccaggcctcc atccagaatg gcaaagacag cacagcccc cagcgtgtcc 1440
 ctgttgctc cccatccgc cacaacatca gcagcagtgg tggagcccca gaccgaacta 1500
 acttcccccg ggggtgtgtcc agccgaagca ccttccatgc tgggcagctc cgacagggtg 1560
 gggaccagca gaatttgccc tacgggtgtga cccagcctc tccctctggc cacagccagg 1620
 gccggcgggg ggcctctggg agcatcttca gcaagttcac ctccaagttt gtacgcagga 1680
 acctgaatga acctgaaagc aaagaccgag tggagacgct cagacctcac gtggtgggca 1740
 gtggcggcaa cgacaaagaa aaggaagaat ttcgggaggc caagccccgc tccctccgct 1800
 tcacgtggag tatgaagacc acgagctcca tggagcccaa cgagatgatg cgggagatcc 1860
 gcaagggtgct ggacgcgaac agctgccaga gcgagctgca tgagaagtac atgctgctgt 1920
 gcatgcacgg cagccggggc cagaggact tcgtgcagtg ggagatggag gtgtgcaaac 1980
 tgccgcggct ctctctcaac ggggttcgat ttaagcggat atcgggcacc tccatggcct 2040
 tcaaaaacat tgcctccaaa atagccaacg agctgaagct ttaaca 2086

<210> 5
 <211> 2248
 <212> DNA
 <213> Homo sapiens

<400> 5
 agtccaacat gattcggggc cgcaactcag ccacctctgc tgatgagcag cccacattg 60
 gaaactaccg gctcctcaag accattggca agggtaattt tgccaagggtg aagttggccc 120
 gacacatcct gactgggaaa gaggtagctg tgaagatcat tgacaagact caactgaact 180

cctccagcct ccagaaacta ttccgcgaag taagaataat gaaggttttg aatcatccca	240
acatagttaa attatttgaa gtgattgaga ctgagaaaac gctctacctt gtcattggagt	300
acgctagtgg cggagaggta tttgattacc tagtggctca tggcaggatg aaagaaaaag	360
aggctcgagc caaattccgc cagatagtgt ctgctgtgca gtactgtcac cagaagttta	420
ttgtccatag agacttaaag gcagaaaacc tgctcttggga tgctgatatg aacatcaaga	480
ttgcagactt tggcttcagc aatgaattca cctttgggaa caagctggac accttctgtg	540
gcagtcccc ttatgctgcc ccagaactct tccagggcaa aaaatatgat ggacccgagg	600
tggatgtgtg gaggctagga gttatcctct atacactggg cagcggatcc ctgccttttg	660
atggacagaa cctcaaggag ctgcgggaac gggctactgag gggaaaatac cgtattccat	720
tctacatgtc cacggactgt gaaaacctgc ttaagaaatt tctcattctt aatcccagca	780
agagaggcac tttagagcaa atcatgaaag atcgatggat gaatgtgggt cacgaagatg	840
atgaactaaa gccttacgtg gagccactcc ctgactacaa ggacccccgg cggacagagc	900
tgatgggtgc catgggttat acacgggaag agatccagga ctgctgggtg ggccagagat	960
acaacgagggt gatggccacc tatctgtctc tgggctacaa gagctccgag ctggaaggcg	1020
acaccatcac cctgaaaccc cggccttcag ctgatctgac caatagcagc gccccatccc	1080
catcccacaa ggtacagcgc agcgtgtcgg ccaatcccaa gcagcggcgc ttcagcgacc	1140
aggctgggtc tgccattccc acctctaatt ctactctaa gaagactcag agtaacaacg	1200
cagaaaataa gcggcctgag gaggaccggg agtcagggcg gaaagccagc agcacagcca	1260
aggtgcctgc cagccccctg cccgggtctgg agaggaagaa gaccacccca accccctcca	1320
cgaacagcgt cctctccacc agcacaatc gaagcaggaa ttccccactt ttggagcggg	1380
ccagcctcgg ccaggcctcc atccagaatg gcaaagacag cctaaccatg ccagggtccc	1440
gggcctccac ggcttctgct tctgccgcag tctctgcggc cgggccccgc cagcaccaga	1500
aatccatgtc ggctccgtg caccocaaca aggcctctgg gctgcccccc acggagagta	1560
actgtgaggt gccgcggccc agcacagccc cccagcgtgt cctgttgcc tccccatccg	1620
cccacaacat cagcagcagt ggtggagccc cagaccgaac taacttcccc cggggtgtgt	1680
ccagccgaag caccttccat gctgggcagc tccgacaggt gggggaccag cagaatttgc	1740
cctacggtgt gacccagcc tctccctctg gccacagcca gggccggcgg ggggcctctg	1800
ggagcatctt cagcaagtcc acctccaagt ttgtacgcag gaacctgaat gaacctgaaa	1860
gcaaagaccg agtggagacg ctgagacctc acgtgggtggg cagtggcggc aacgacaaag	1920
aaaaggaaga atttcgggag gccaaagccc gctccctccg cttcacgtgg agtatgaaga	1980
ccacgagctc catggagccc aacgagatga tgcgggagat ccgcaagggt ctggacgcga	2040

acagctgccca gagcgagctg catgagaagt acatgctgct gtgcatgcac ggcacgcccgg 2100
gccacgagga cttcgtgcag tgggagatgg aggtgtgcaa actgccgcccgt ctctctctca 2160
acgggggttcg atttaagcgg atatcgggca cctccatggc cttcaaaaac attgcctcca 2220
aaatagccaa cgagctgaag ctttaaca 2248

<210> 6
<211> 2701
<212> DNA
<213> Homo sapiens

<400> 6
ggcacgaggg ctgaacgaga gggacacgga gcagcccacc ttgggacacc ttgactccaa 60
gcccagcagt aagtccaaca tgattcgggg cgcgaactca gccacctctg ctgatgagca 120
gccccacatt ggaaactacc ggctcctcaa gaccattggc aagggttaatt ttgccaaggt 180
gaagttggcc cgacacatcc tgactgggaa agaggtagct gtgaagatca ttgacaagac 240
tcaactgaac tcctccagcc tccagaaact attccgcgaa gtaagaataa tgaaggtttt 300
gaatcatccc aacatagtta aattatattga agtgattgag actgagaaaa cgctctacct 360
tgtcatggag tacgctagtg gcggagaggt atttgattac ctagtggctc atggcaggat 420
gaaagaaaaa gaggctcgag ccaaattccg ccagatagtg tctgctgtgc agtactgtca 480
ccagaagttt attgtccata gagacttaaa ggcagaaaac ctgctccttg atgctgatat 540
gaacatcaag attgcagact ttggcttcag caatgaattc acctttggga acaagctgga 600
caccttctgt ggcagtcacc cttatgctgc ccagaaactc ttccagggca aaaaatatga 660
tggacccgag gtggatgtgt ggagcctagg agttatcctc tatacactgg tcagcggatc 720
cctgcctttt gatggacaga acctcaagga gctgcgggaa cgggtactga ggggaaaata 780
ccgtattcca ttctacatgt ccacggactg tgaaaacctg cttaagaaat ttctcattct 840
taatcccagc aagagaggca ctttagagca aatcatgaaa gatcgatgga tgaatgtggg 900
tcacgaagat gatgaactaa agccttacgt ggagccactc cctgactaca aggacccccg 960
gcggacagag ctgatgggtg ccatgggtta tacacgggaa gagatccagg actcgctggt 1020
gggccagaga tacaacgagg tgatggccac ctatctgctc ctgggctaca agagctccga 1080
gctggaaggc gacaccatca ccctgaaacc ccggccttca gctgatctga ccaatagcag 1140
cgccccatcc ccatcccaca aggtacagcg cagcgtgtcg gccaatccca agcagcggcg 1200
cttcagcgac caggcagctg gtcctgccat tcccacctct aattcttact ctaagaagac 1260
tcagagtaac aacgcagaaa ataagcggcc tgaggaggac cgggagtcag ggcggaagc 1320
cagcagcaca gccaaaggtg ctgccagccc cctgcccggg ctggagagga agaagaccac 1380

```

ccccaccccc tccacgaaca gcgtcctctc caccagcaca aatcgaagca ggaattcccc 1440
acttttggag cgggccagcc tcggtcaggg ctccatccag aatggcaaag acagcctaac 1500
catgccaggg tcccgggcct ccacggcttc tgcttctgcc gcagtctctg cggcccggcc 1560
ccgccagcac cagaaatcca tgtcggcctc cgtgcacccc aacaaggcct ctgggctgcc 1620
ccccacggag agtaactgtg aggtgccgcg gccacgcaca gccccccagc gtgtccctgt 1680
tgcttccccca tccgcccaaca acatcagcag cagtgggtgga gcccagacc gaactaactt 1740
cccccggggt gtgtccagcc gaagcacctt ccatgctggg cagctccgac aggtgctggga 1800
ccagcagaat ttgccctacg gtgtgacccc agcctctccc tctggccaca gccagggccg 1860
gcgggggggc tctgggagca tcttcagcaa gttcacctcc aagtttgtac gcagaaatct 1920
gtctttcagg tttgccagaa ggaacctgaa tgaacctgaa agcaaagacc gagtggagac 1980
gctcagacct cacgtggtgg gcagtggcgg caacgacaaa gaaaaggaag aatttcggga 2040
ggccaagccc cgtccctcc gcttcacgtg gagtatgaag accacgagct ccatggagcc 2100
caacgagatg atgcgggaga tccgcaaggt gctggacgcg aacagctgcc agagcgagct 2160
gcatgagaag tacatgctgc tgtgcatgca cggcacgccc ggccacgagg acttcgtgca 2220
gtgggagatg gaggtgtgca aactgccgcg gctctctctc aacgggggttc gatttaagcg 2280
gatatcgggc acctccatgg ccttcaaaaa cattgcctcc aaaatagcca acgagctgaa 2340
gctttaacag gctgccagga gcggggggcg cggggggcggg ccagctggac gggctgccgg 2400
ccgctgcgcc gccccacctg ggcgagactg cagcgatgga ttggtgtgtc tcccctgctg 2460
gcacttctcc cctccctggc ccttctcagt tttctcttac atgtttgtgg ggggtgggag 2520
attgttctcc agccccccac attcaccct gccacagatg tccccctct cctctccct 2580
actggaggca aaggaagggg aggggtggatg ggggggcagg gctccccctc ggtactgcgg 2640
ttgcacagag tatttcgcct aaaccaagaa attttttatt accaaaaaaaa aaaaaaaaaa 2700
a 2701

```

<210> 7

<211> 2112

<212> DNA

<213> Homo sapiens

<400> 7

```

cccagcagta agtccaacat gattcggggc cgcaactcag ccacctctgc tgatgagcag 60
ccccacattg gaaactaccg gctcctcaag accattggca agggtaattt tgccaagggtg 120
aagttggccc gacacatcct gactgggaaa gaggtagctg tgaagatcat tgacaagact 180
caactgaact cctccagcct ccagaaacta ttccgcgaag taagaataat gaaggttttg 240
aatcatccca acatagttaa attatttgaa gtgattgaga ctgagaaaac gctctacctt 300

```

```

gtcatggagt acgctagtgg cggagaggta tttgattacc tagtggctca tggcaggatg      360
aaagaaaaag aggctcgagc caaattccgc cagatagtgt ctgctgtgca gtactgtcac      420
cagaagttta ttgtccatag agacttaaag gcagaaaacc tgctcttgga tgctgatatg      480
aacatcaaga ttgcagactt tggcttcagc aatgaattca cctttgggaa caagctggac      540
accttctgtg gcagtcccc ttatgctgcc ccagaactct tccagggcaa aaaatatgat      600
ggacccgagg tggatgtgtg gagcctagga gttatcctct atacactggg cagcggatcc      660
ctgccttttt atggacagaa cctcaaggag ctgcgggaac gggactgag gggaaaatac      720
cgtattccat tctacatgtc cacggactgt gaaaacctgc ttaagaaatt tctcattctt      780
aatcccagca agagaggcac tttagagcaa atcatgaaag atcgatggat gaatgtgggt      840
cacgaagatg atgaactaaa gccttacgtg gagccactcc ctgactacaa ggacccccgg      900
cggacagagc tgatgggtgc catgggttat acacgggaag agatccagga ctgcgtgggtg      960
ggccagagat acaacgaggt gatggccacc tatctgctcc tgggctacaa gagctccgag     1020
ctggaaggcg acaccatcac cctgaaacct cggccttcag ctgatctgac caatagcagc     1080
gccccatccc catcccacaa ggtacagcgc agcgtgtcgg ccaatcccaa gcagcggcgc     1140
ttcagcgacc aggctgggtc tgccattccc acctctaatt cttactctaa gaagactcag     1200
agtaacaacg cagaaaataa gcggcctgag gaggaccggg agtcagggcg gaaagccagc     1260
agcacagcca aggtgcctgc cagccccctg cccgggtctgg agaggaagaa gaccaccca     1320
acccccctca cgaacagcgt cctctccacc agcacaatc gaagcaggaa ttccccactt     1380
ttggagcggg ccagcctcgg ccaggcctcc atccagaatg gcaaagacag cacagcccc     1440
cagcgtgtcc ctgttgctc cccatccgcc cacaacatca gcagcagtgg tggagcccca     1500
gaccgaacta acttcccccg ggggtgtgtc agccgaagca ccttccatgc tgggcagctc     1560
cgacaggtgc gggaccagca gaatttgccc tacgggtgtga cccagcctc tccctctggc     1620
cacagccagg gccggcgggg ggctctggg agcatcttca gcaagttcac ctccaagttt     1680
gtacgcagga acctgaatga acctgaaagc aaagaccgag tggagacgct cagacctcac     1740
gtggtgggca gtggcggcaa cgacaaagaa aaggaagaat ttcgggaggc caagccccgc     1800
tccctccgct tcacgtggag tatgaagacc acgagctcca tggagcccaa cgagatgatg     1860
cgggagatcc gcaaggtgct ggacgcgaac agctgccaga gcgagctgca tgagaagtac     1920
atgctgctgt gcatgcacgg cacgccgggc cacgaggact tcgtgcagtg ggagatggag     1980
gtgtgcaaac tgccgcggct ctctctcaac ggggttcgat ttaagcggat atcgggcacc     2040
tccatggcct tcaaaaacat tgcctccaaa atagccaacg agctgaagct ttaacaggct     2100
gccaggagcg gg                                         2112

```

<210> 8
 <211> 2965
 <212> DNA
 <213> Homo sapiens

<400> 8
 cgggcaaccg cctcgcccgga agccctccct cgttactgtc cgcatacccc ggcgggcgccg 60
 ccgcgggaaag cgggtccccc tcctcttact ccgcgtcctc ttccctcttt cccccgcggg 120
 ggcacgcttg ttgcaccgtc ccgcggcctg cgggagccgc tcgccccgga cttgagctcg 180
 cgtacgaccc atttcctgtc gccccccgga gcccgcacca cagcccggcc ggtctagacc 240
 ccggcagacc ccgctggccg cacaaaatgt cggcccggac gccattgccg acggtgaacg 300
 agcgggacac ggtaaatcat acgactgtgg atggatatac tgaaccacac atccagccta 360
 ccaagtgcag tagcagacag aacatcccc ggtgtagaaa ctccattacg tcagcaacag 420
 atgaacagcc tcacattgga aattacogtt tacaaaaaac aatagggaag ggaaattttg 480
 ccaaagtcaa attggcaaga cacgttctaa ctggtagaga ggttgctgtg aaaataatag 540
 acaaaaactca gctaaatcct accagtctac aaaagttatt tcgagaagta cgaataatga 600
 agatactgaa tcatccta atagtaaaat tgtttgaagt tattgaaaca gagaagactc 660
 tctatttagt catggaatac gcgagtgggg gtgaagtatt tgattactta gttgcccatg 720
 gaagaatgaa agagaaagag gcccgtgcaa aatttaggca gattgtatct gctgtacagt 780
 attgtcatca aaagtacatt gttcacctgt atcttaaggc tgaaaacctt ctcccttgatg 840
 gtgatatgaa tattaaaatt gctgactttg gtttttagtaa tgaatttaca gttgggaaca 900
 aattggacac attttgtgga agcccaccct atgctgctcc cgagcttttc caaggaaaga 960
 agtatgatgg gcctgaagtg gatgtgtgga gtctgggcgt cattctctat acattagtca 1020
 gtggctcctt gcctttcgat ggccagaatt taaaggaact gcgagagcga gttttacgag 1080
 ggaagtaccg tattcccttc tatatgtcca cagactgtga aaatcttctg aagaaattat 1140
 tagtcctgaa tccaataaag agaggcagct tggaacaaat aatgaaagat cgatggatga 1200
 atgttggtca tgaagaggaa gaactaaagc catatactga gcctgatccg gatttcaatg 1260
 acacaaaaag aatagacatt atggtcacca tgggctttgc acgagatgaa ataaatgatg 1320
 ccttaataaaa tcagaagtat gatgaagtta tggctactta tattcttcta ggtagaaaac 1380
 cacctgaatt tgaagggtgt gaatcgttat ccagtggaaa cttgtgtcag aggtcccgcc 1440
 ccagtagtga cttaaacaac agcactcttc agtcccctgc tcacctgaag gtccagagaa 1500
 gtatctcagc aaatcagaag cagcggcggt tcagtgatca tgctggtcca tccattcctc 1560
 ctgctgtatc atataccaaa agacctcagg ctaacagtgt ggaaagtga cagaaagagg 1620

agtgggacaa agatgtggct cgaaaacttg gcagcacaaac agttggatca aaaagcgaga 1680
 tgactgcaag ccctcttgta gggccagaga ggaaaaaatc ttcaactatt ccaagtaaca 1740
 atgtgtattc tggaggtagc atggcaagaa ggaatacata tgtctgtgaa aggaccacag 1800
 atcgatacgt agcattgcag aatggaaaag acagcagcct tacggagatg tctgtgagta 1860
 gcatactctt tgcaggctct tctgtggcct ctgctgtccc ctcagcacga ccccgccacc 1920
 agaagtccat gtccacttct ggtcatccta ttaaagtcac actgccaaacc attaaagacg 1980
 gctctgaagc ttaccggcct ggtacaaccc agagagtgcc tgctgtctcc ccactgtctc 2040
 acagtattag tactgcgact ccagaccgga cccgttttcc ccgagggagc tcaagccgaa 2100
 gcactttcca tgggtgaacag ctccgggagc gacgcagcgt tgcttataat gggccacctg 2160
 cttcaccatc ccatgaaacg ggtgcatttg cacatgccag aaggggaacg tcaactggta 2220
 taataagcaa aatcacatcc aaatttgctc gcagggatcc aagtgaaggc gaagccagtg 2280
 gcagaaccga cacctcaaga agtacatcag gggaaaccaa agaaagagac aaggaagagg 2340
 gtaaagattc taagccgcgt tctttgcggt tcacatggag tatgaagacc actagttcaa 2400
 tggaccctaa tgacatgatg agagaaatcc gaaaagtgtt agatgcaaat aactgtgatt 2460
 atgagcaaaa agagagattt ttgcttttct gtgtccatgg agacgctaga caggatagcc 2520
 tcgtgcagtg ggagatggaa gtctgcaagt tgccacgact gtcacttaat ggggttcgct 2580
 tcaagcgaat atctgggaca tctattgcct ttaagaacat tgcataaaaa atagcaaatg 2640
 agcttaagct gtaaagaagt ccaaatttac aggttcaggg aagatacata catatatgag 2700
 gtacagtttt tgaatgtact ggtaatgcct aatgtggtct gcctgtgaat ctccccatgt 2760
 agaatttgcc cttaatgcaa taaggttata catagtattg aactgtaaaa ttaaagtcag 2820
 tatgaactat aataaatatc tgtagcttaa aaagtaggtt cacatgtaca ggtaagtata 2880
 ttgtgtattt ctgttcattt tctgttcata gagttgtata ataaaacatg attgcttaaa 2940
 aacttgaaaa aaaaaaaaaa aaaaa 2965

<210> 9
 <211> 3210
 <212> DNA
 <213> Homo sapiens

<400> 9
 ggcgcggcgg cggcggtggc tgtgaccgcg cggaccgagc cgagacattc gcgcgggggg 60
 atcgggcgcc gccgccgtg ggcgccgggc gcgtggatgc ggctgggtcg ggcggcgccg 120
 tacacctgag gcggagaacg ggcgcggcg cgggtgacgc tgtcagggcc gcggttcctg 180
 acgcccaggc gctcgccagg acgagccagg cagtgatttg aggcaccggc ttcaccttca 240
 cccatggtcc ggagagccta gcggggctcg ccaccgcctc ccggtcctcc ttccacgctc 300

catcctgccca gcctcgccgc ccgcgcagcg ccgggcaacc gcctcgcccg aagccctccc	360
tcgttactgt ccgcataccc cggcggcgcc gccgcgggaa ggggctcccc ctccctcttc	420
tccgcgtcct ctccctctt tccccgcgc gggcgcgttg ttgcaccgce ccgcggcctg	480
cgggagccgc tcgccccggc cttgtgctcg cgtccgcacc cctttcctgt cggcccccg	540
ggccccgcacc acagcccggc cggcgagacc ccggccagac cccgctgccc gcacaaaatg	600
tcggccccga cgccattgcc gacggtgaac gagcgggaca cggaaaatca tacatctgtg	660
gatggatata ctgaaccaca catccagcct accaagtcga gtagcagaca gaacatcccc	720
cgggtgtagaa actccattac gtcagcaaca gatgaacagc ctacattgg aaattaccgt	780
ttacaaaaaa caatagggaa gggaaatttt gccaaagtca aattggcaag acacgttcta	840
actggtagag aggttgctgt gaaaataata gacaaaactc agctaaatcc taccagtcta	900
caaaagttat ttcgagaagt acgaataatg aagatactga atcatcctaa tatagggtgaa	960
gtatttgatt acttagttgc ccatggaaga atgaaagaga aagaggcccg tgcaaaattt	1020
aggcagattg tatctgctgt acagtattgt catcaaaagt acattgttca ccgtgatctt	1080
aagctgaaaa ccttctcctt gatggtgata tgaatattaa aattgctgac tttggtttta	1140
gtaatgaatt tacagttggg aacaaattgg acacattttg tggaagccca ccctatgctg	1200
ctcccagact tttccaagga aagaagtatg atgggcctga agtggatgtg tggagtctgg	1260
gcgtcattct ctatacatta gtcagtggct ccttgccttt cgatggccag aatttaaagg	1320
aactgcgaga gcgagtttta cgagggaagt accgtattcc cttctatatg tccacagact	1380
gtgaaaatct tctgaagaaa ttattagtcc tgaatccaat aaagagaggc agcttggaac	1440
aaataatgaa agatcgatgg atgaatgttg gtcatgaaga ggaagaacta aagccatata	1500
ctgagcctga tccggatttc aatgacacaa aaagaataga cattatggtc accatgggct	1560
ttgcacgaga tgaaataaat gatgccttaa taaatcagaa gtatgatgaa gttatggcta	1620
cttatattct tctaggtaga aaaccacctg aatttgaagg tggatgaatcg ttatccagt	1680
gaaacttggtg tcagaggtcc cggcccagta gtgacttaaa caacagcact cttcagtccc	1740
ctgctcacct gaagggtccag agaagtatct cagcaaatca gaagcagcgg cgtttcagt	1800
atcatgctgg tccatccatt cctcctgctg tatcatatac caaaagacct caggctaaca	1860
gtgtggaaag tgaacagaaa gaggagtggg acaaagatgt ggctcgaaaa cttggcagca	1920
caacagttgg atcaaaaagc gagatgactg caagccctct tgtagggcca gagaggaaaa	1980
aatcttcaac tattccaagt aacaatgtgt attctggagg tagcatggca agaaggaata	2040
catatgtctg tgaaaggacc acagatcgat acgtagcatt gcagaatgga aaagacagca	2100
gccttacgga gatgtctgtg agtagcatat cttctgcagg ctcttctgtg gcctctgctg	2160

tccccctcagc acgacccccgc caccagaagt ccatgtccac ttctgggtcat cctattaaag 2220
 tcacactgcc aaccattaaa gacggctctg aagcttaccg gcctgggtaca acccagagag 2280
 tgccctgctgc ttccccatct gctcacagta ttagtactgc gactccagac cggacccggt 2340
 ttccccgagg gagctcaagc cgaagcactt tccatgggtga acagctccgg gagcgacgca 2400
 gcgttgctta taatggggcca cctgcttcac catcccatga aacgggtgca tttgcacatg 2460
 ccagaagggg aacgtcaact ggtataataa gcaaaatcac atccaaattt gttcgcaggg 2520
 atccaagtga aggcgaagcc agtggcagaa ccgacacctc aagaagtaca tcaggggaac 2580
 caaaagaaag agacaaggaa gagggtaaag attctaagcc gcgttccttg cggttcacat 2640
 ggagtatgaa gaccactagt tcaatggacc ctaatgacat gatgagagaa atccgaaaag 2700
 tgttagatgc aaataactgt gattatgagc aaaaagagag atttttgctt ttctgtgtcc 2760
 atggagacgc tagacaggat agcctcgtgc agtgggagat ggaagtctgc aagttgccac 2820
 gactgtcact taatgggggtt cgcttcaagc gaatatctgg gacatctatt gcctttaaga 2880
 acattgcac aaaaatagca aatgagctta agctgtaaag aagtccaaat ttacaggttc 2940
 agggaagata catacatata tgaggtagac tttttgaatg tactggtaat gcctaattgtg 3000
 gtctgcctgt gaatctcccc atgtagaatt tgcccttaat gcaataaggt tatacatagt 3060
 tatgaactgt aaaattaaag tcagtatgaa ctataataaa tatctgtagc ttaaaaagta 3120
 gggttcacatg tacaggtaag tatattgtgt atttctgttc attttctgtt catagagtgt 3180
 tataataaaa catgattgct taaaaacttg 3210

<210> 10

<211> 2505

<212> DNA

<213> Homo sapiens

<400> 10

gctggccgca caaaatgtcg gcccggaagc cattgccgac ggtgaacgag cgggacacgg 60
 aaaatcatac atctgtggat ggatatactg aaccacacat ccagcctacc aagtcgagta 120
 gcagacagaa catcccccg tgtagaaact ccattacgtc agcaacagat gaacagcctc 180
 acattggaaa ttaccgttta caaaaaacaa tagggaaggg aaattttgcc aaagtcaaat 240
 tggcaagaca cgttctaact ggtagagagg ttgctgtgaa aataatagac aaaactcagc 300
 taaatcctac cagtctacaa aagttatttc gagaagtacg aataatgaag atactgaatc 360
 atcctaatat agtaaaattg tttgaagtta ttgaaacaga gaagactctc tatttagtca 420
 tggaaacgc gagtgggggt gaagtatttg attacttagt tgcccatgga agaataagaa 480
 agaaagaggc ccgtgcaaaa ttaggcaga ttgtatctgc tgtacagtat tgtcatcaaa 540

agtacattgt	tcaccgtgat	cttaaggctg	aaaaccttct	ccttgatggt	gatatgaata	600
ttaaaattgc	tgacttttgg	tttagtaatg	aatttacagt	tgggaacaaa	ttggacacat	660
tttgtggaag	cccaccctat	gctgctcccg	agcttttcca	aggaaagaag	tatgatgggc	720
ctgaagtgga	tgtgtggagt	ctgggcgtca	ttctctatac	attagtcagt	ggctccttgc	780
ctttcgatgg	ccagaattta	aaggaaactgc	gagagcgagt	tttacgaggg	aagtaccgta	840
ttcccttcta	tatgtccaca	gactgtgaaa	atcttctgaa	gaaattatta	gtcctgaatc	900
caataaagag	aggcagcttg	gaacaaataa	tgaaagatcg	atggatgaat	gttggtcatg	960
aagaggaaga	actaaagcca	tatactgagc	ctgatccgga	tttcaatgac	acaaaaagaa	1020
tagacattat	ggtcaccatg	ggctttgcac	gagatgaaat	aaatgatgcc	ttaataaatc	1080
agaagtatga	tgaagttaatg	gctacttata	ttcttctagg	tagaaaacca	cctgaatttg	1140
aaggtggtga	atcgttatcc	agtggaaact	tgtgtcagag	gtcccggccc	agtagtgact	1200
taaacaacag	cactcttcag	tcccctgctc	acctgaaggt	ccagagaagt	atctcagcaa	1260
atcagaagca	gcggcgtttc	agtgatcatg	ctgggtccatc	cattcctcct	gctgtatcat	1320
ataccaaaag	acctcaggct	aacagtgtgg	aaagtgaaca	gaaagaggag	tgggacaaag	1380
atgtggctcg	aaaacttggc	agcacaacag	ttggatcaaa	aagcgagatg	actgcaagcc	1440
ctcttgtagg	gccagagagg	aaaaaatctt	caactattcc	aagtaacaat	gtgtattctg	1500
gaggtagcat	ggcaagaagg	aatacatatg	tctgtgaaag	gaccacagat	cgatacgtag	1560
cattgcagaa	tggaaaagac	agcagcctta	cggagatgtc	tgtgagtagc	atatcttctg	1620
caggctcttc	tgtggcctct	gctgtcccct	cagcacgacc	ccgccaccag	aagtccatgt	1680
ccacttctgg	tcatcctatt	aaagtcacac	tgccaaccat	taaagacggc	tctgaagctt	1740
accggcctgg	tacaaccag	agagtgcctg	ctgcttcccc	atctgctcac	agtattagta	1800
ctgcgactcc	agaccggacc	cgttttcccc	gagggagctc	aagccgaagc	actttccatg	1860
gtgaacagct	ccgggagcga	cgcagcgttg	cttataatgg	gccacctgct	tcaccatccc	1920
atgaaacggg	tgcatttgca	catgccagaa	ggggaacgtc	aactggtata	ataagcaaaa	1980
tcacatccaa	atttgttcgc	agggatccaa	gtgaaggcga	agccagtggc	agaaccgaca	2040
cctcaagaag	tacatcaggg	gaacaaaaag	aaagagacaa	ggaagagggg	aaagattcta	2100
agccgcgttc	tttgoggttc	acatggagta	tgaagaccac	tagttcaatg	gaccctaata	2160
acatgatgag	agaaatccga	aaagtgttag	atgcaaataa	ctgtgattat	gagcaaaaag	2220
agagattttt	gcttttctgt	gtccatggag	acgctagaca	ggatagcctc	gtgcagtggg	2280
agatggaagt	ctgcaagtgg	ccacgactgt	cacttaaatg	ggttcgcttc	aagcgaatat	2340
ctgggacatc	tattgccttt	aagaacattg	catcaaaaat	agcaaatgag	cttaagctgt	2400

aaagaagtcc aaatttacag gttcagggaa gatacatata tatatgaggt acagttttttg 2460
 aatgtactgg taatgcctaa tgtggtctgc ctgtgaatct cccca 2505

<210> 11
 <211> 4638
 <212> DNA
 <213> Homo sapiens

<400> 11
 ggcgcggcgg cggcgggtggc tgtgaccgcg cggaccgagc cgagacattc gcgccggggg 60
 atcggggcgcc gccgcgcgtg ggccccgggc gcgtggatgc ggctgggtcg ggccggcgccg 120
 tacacctgag gcggagaacg gggcgcgggcg cgggtgacgc tgtcagggcc gcggttcctg 180
 acgcccaggc gctcgccagg acgagccagg cagtgatttg aggcaccggc ttcaccttca 240
 cccatgggtcc ggagagccta gcgggggctcg ccaccgcctc ccggctcccc ttccacgcct 300
 catcctgccca gcctcgccgc cccgccagcg ccgggcaacc gcctcgcccg aagccctccc 360
 tcgttactgt ccgcataccc cggcggcgcc gccgcgggaa gcggctcccc ctctcttccc 420
 tccgcgtcct ctctcctctt tccccgcgcg gggcgcgttg ttgcaccgcc ccgcggcctg 480
 cgggagccgc tcgccccggc cttgtgctcg cgtccgcacc ctttctctgt cgcggcccg 540
 ggccccgacc acagcccgcc cggcgagacc ccggccagac cccgctgccc gcacaaaatg 600
 tcggccccga cgccattgcc gacgggtgaac gagcgggaca cggaaaatca tacatctgtg 660
 gatggatata ctgaaccaca catccagcct accaagtcca gtagcagaca gaacatcccc 720
 cgggtgtagaa actccattac gtcagcaaca gatgaacagc ctacattgg aaattaccgt 780
 ttacaaaaaa caatagggaa gggaaatatt gccaaagtca aattggcaag acacgttcta 840
 actggtagag aggttgctgt gaaaataata gacaaaactc agctaaatcc taccagtcta 900
 caaaagtatt ttcgagaagt acgaataatg aagatactga atcatcctaa tatagggtgaa 960
 gtatttgatt acttagttgc ccatggaaga atgaaagaga aagaggcccg tgcaaaatth 1020
 aggcagattg tatctgctgt acagtattgt catcaaaagt acattgttca ccgtgatctt 1080
 aaggctgaaa accttctcct tgatgggtgat atgaatatta aaattgctga ctttggtttt 1140
 agtaatgaat ttacagttgg gaacaaattg gacacatttt gtggaagccc accctatgct 1200
 gctcccgagc ttttccaagg aaagaagtat gatgggcctg aagtggatgt gtggagtctg 1260
 ggcgtcattc tctatacatt agtcagtggc tccttgccct tcgatggcca gaatttaaag 1320
 gaactgcgag agcgagtttt acgagggaaag taccgtattc ccttctatat gtccacagac 1380
 tgtgaaaatc ttctgaagaa attattagtc ctgaatccaa taaagagagg cagcttgga 1440
 caaataatga aagatcgatg gatgaatggt ggtcatgaag aggaagaact aaagccatat 1500
 actgagcctg atccggattt caatgacaca aaaagaatag acattatggt caccatgggc 1560

tttgcacgag atgaaataaa tgatgcctta ataaatcaga agtatgatga agttatggct 1620
 acttatattc ttctaggtag aaaaccacct gaatttgaag gtggtgaatc gttatccagt 1680
 ggaaacttgt gtcagaggtc cgggccagct agtgacttaa acaacagcac tcttcagctc 1740
 cctgctcacc tgaaggcca gagaagtatc tcagcaaatc agaagcagcg gcgtttcagt 1800
 gatcatgctg gtccatccat tctcctgct gtatcatata ccaaagacc tcaggctaac 1860
 agtgtggaag gtgaacagaa agaggagtgg gacaaagatg tggctcgaaa acttggcagc 1920
 acaacagttg gatcaaaaag cgagatgact gcaagcctc ttgtagggcc agagaggaaa 1980
 aaatcttcaa ctattccaag taacaatgtg tattctggag gtagcatggc aagaaggaat 2040
 acatatgtct gtgaaaggac cacagatcga tacgtagcat tgcagaatgg aaaagacagc 2100
 agccttacgg agatgtctgt gagtagcata tcttctgcag gctcttctgt ggcctctgct 2160
 gtccctcag cagcaccctg ccaccagaag tccatgtcca cttctggcca tcctattaaa 2220
 gtcacactgc caaccattaa agacggctct gaagcttacc ggcctggtag aaccagaga 2280
 gtgcctgctg cttcccatc tgctcacagt attagtactg cgactccaga ccggaccctg 2340
 tttcccgag ggagctcaag ccgaagcact tccatgggtg aacagctccg ggagcgacgc 2400
 agcgttgctt ataatgggct acctgcttca ccatcccatg aaacgggtgc atttgacat 2460
 gccagaaggg gaacgtcaac tgggtataata agcaaatca catccaaatt tgttcgaga 2520
 agtacatcag gggaacaaaa agaaagagac aaggaagagg gtaaagattc taagccgcgt 2580
 tctttgcggt tcacatggag tatgaagacc actagttaa tggaccctaa tgacatgatg 2640
 agagaaatcc gaaaagtgtt agatgcaaat aactgtgatt atgagcaaaa agagagattt 2700
 ttgcttttct gtgtccatgg agacgctaga caggatagcc tcgtgcagtg ggagatggaa 2760
 gtctgcaagt tgccacgact gtcacttaat ggggttcgct tcaagcgaat atctgggaca 2820
 tctattgcct ttaagaacat tgcataaaaa atagcaaatg agcttatgct gtaaagaagt 2880
 ccaaatttac aggttcaggg aagatacata catatatgag gtacagtttt tgaatgtact 2940
 ggtaatgcct aatgtggtct gcctgtgaat ctcccatgt agaatttgcc cttaatgcaa 3000
 taaggttata catagtattg aactgtaaaa ttaaagtcag tatgaactat aataaatatc 3060
 tgtagcttaa aaagtaggtt cacatgtaca ggtaagtata ttgtgtattt ctgttcattt 3120
 tctgttcata gagttgtata ataaaacatg attgctttaa aacttgtata gttgtctaga 3180
 tttctgcacc tgaatgtatg ttgatgctt tgatttgaaa atgttcttcc ctgttattta 3240
 cattctgggtg ggtttttaa attcttacct ccatcatgca attttgaaaa ttgtgtccag 3300
 aattaaaagt gcatagaaat agcctttaca attgtagcat ggacctttaa aaattgtttt 3360
 aaaatcttat ttaaatttaa accagaagct gaaaaataga tcagctttat tatacacaaa 3420

```

attattactg cttatctttg ctcttttcoct tgttatcccg caaggtttag ttgagaagat 3480
acaaaatggt tacagtgttg gcacttagag tttttaaatt caagtacatg aaattcagta 3540
atagcattgc cttgagctaa ctaggaagta cccgggaaaaa agttaaatct acatcaagtt 3600
tcttttgaac tttgaagtgt tttctgaccc actgctaact gtagcaacaa aattttaaag 3660
aaaaaaaaaca tactttatct ggctattata acataaactg tcacgtaggt ttgctgcctt 3720
cagaataccg caatttaatt gcgggaatat aataatattg ggactgtttc acagcacaaa 3780
ctcatcttta cagtgttgat caatgcatca gttaagaaat aatgccacct caggaattaa 3840
ctggcattgg gaacatttgc ctcatctccc tgctatcctc ttcattcacc cctgccactg 3900
taatatttat aagtacttaa gagacttgtg agcaaacat actatttata acagtatatg 3960
attgatttat gcttatgtgg ttgttcagtt tgttcccatg taactcgttt gttttaaata 4020
ttttgccaga tttcttgtat ttattccaca tcattatgcc tataatgtgc cgctttgtga 4080
ttgggcattt gctactttt ctttcataat tagtgatata tgcgatgtaa aaccactagt 4140
aaaggtagat ttttaatactt gttattttat actgaattag ccttggaggt tgactgtgca 4200
atgttattta ctgttgtaat tactgtaata ccaacatag gcccccatct gcacactcct 4260
gaaaaacaga aagtgtattc aaattttatc agtttaaaga aaataaagct gtgataaata 4320
ctgtaattcc aacctacatt agaagggtcta agtgtaggtg atgtgccatt ccataatggc 4380
ttccagacta ggggtgaattt tatgttctgt actgtactgt gatgtagctt tcttctgtaa 4440
cagttatggt ttaaaattaa gtgagttttt tttttgcctt agcaaagggt ggtgtttgaa 4500
aaaaaaaaatg tgtagccctt ttttaacctg gtgttcattc aaaaaaaaaat tgatgcaaat 4560
ctttattcac tttcactggg gcacactgaa attttacttg aacagttctc ataataaagc 4620
acttgtcttt tgctcttt 4638

```

```

<210> 12
<211> 2720
<212> DNA
<213> Homo sapiens

```

```

<400> 12
tcatggaata cgcgagtggg ggtgaagtat ttgattactt agttgcccat ggaagaatga 60
aagagaaaga ggcccggtgca aaatttaggc agattgtatc tgctgtacag tattgtcatc 120
aaaagtacat tgttcaccgt gatcttaagg ctgaaaacct tctccttgat ggtgatatga 180
atattaaaat tgctgacttt ggtttttagta atgaatttac agttgggaac aaattggaca 240
cattttgtgg aagcccaccc tatgtgtctc ccgagctttt ccaaggaaag aagtatgatg 300
gtcctgaagt ggatgtgtgg agtctgggcg tcattctcta tacattagtc agtggctcct 360

```

tgccttttoga	tggccagaat	ttaaaggaac	tgcgagagcg	agtttttacga	gggaagtacc	420
gtattccctt	ctatatgtcc	acagactgtg	aaaatcttct	gaagaaatta	ttagtccctga	480
atccaataaa	gagaggcagc	ttggaacaaa	taatgaaaga	tcgatggatg	aatgttggtc	540
atgaagagga	agaactaaag	ccatatactg	agcctgatcc	ggatttcaat	gacacaaaaa	600
gaatagacat	tatggtcacc	atgggctttg	cacgagatga	aataaatgat	gccttaataa	660
atcagaagta	tgatgaagtt	atggctactt	atattcttct	aggtagaaaa	ccacctgaat	720
ttgaagggtg	tgaatcgta	tccagtggaa	acttgtgtca	gaggtcccg	ccagtagtg	780
acttaaacia	cagcactctt	cagtccctg	ctcacctgaa	ggccagaga	agtatctcag	840
caaatacaga	gcagcggcgt	ttcagtgatc	atgctgggtc	atccattcct	cctgctgtat	900
catataccaa	aagacctcag	gctaacagt	tggaaagtga	acagaaagag	gagtgggaca	960
aagatgtggc	tcgaaaactt	ggcagcacia	cagttggatc	aaaaagcgag	atgactgcaa	1020
gccctcttgt	agggccagag	aggaaaaaat	cttcaactat	tccaagtaac	aatgtgtatt	1080
ctggaggtag	catggcaaga	aggaatacat	atgtctgtga	aaggaccaca	gatcgatacg	1140
tagcattgca	gaatggaaaa	aacagcagcc	ttacggagat	gtctgtgagt	agcatatctt	1200
ctgcaggctc	ttctgtggcc	tctgtgtccc	cctcagcacg	accccgccac	cagaagtcca	1260
tgtccacttc	tggatcatcct	attaaagtca	cactgccaac	cattaaagac	ggctctgaag	1320
cttaccggcc	tggatacaacc	cagagagtgc	ctgtgtcttc	cccatctgct	cacagtatta	1380
gtactgcgac	tccagaccgg	accggttttc	cccaggggag	ctcaagccga	agcactttcc	1440
atggtgaaca	gctccgggag	cgacgcagcg	ttgcttataa	tgggccacct	gcttcaccat	1500
cccatgaaac	gggtgcattt	gcacatgcca	gaaggggaac	gtcaactggt	ataataagca	1560
aaatcacatc	caaatttgtt	cgcagaagta	catcagggga	acaaaaagaa	agagacaagg	1620
aagagggtaa	agattctaag	ccgcgttctt	tgcgggtcac	atggagtatg	aagaccacta	1680
gttcaatgga	ccctaataac	atgatgagag	aaatccgaaa	agtgttagat	gcaaataact	1740
gtgattatga	gcaaaaagag	agatttttgc	ttttctgtgt	ccatggagac	gctagacagg	1800
atagcctcgt	gcagtgggag	atggaagtct	gcaagttgca	cgactgtcac	ttaatggggt	1860
tcgcttcaag	cgaatatctg	ggacatctat	tgcctttaag	aacattgcat	caaaaatagc	1920
aaatgagctt	aagctgtaaa	gaagtccaaa	tttacagggt	caggggaagat	acatacatat	1980
atgaggtaca	gtttttgaat	gtactggtaa	tgcctaattg	ggtctgcctg	tgaatctccc	2040
catgtagaat	ttgcccttaa	tgcaataagg	ttatacatag	ttatgaactg	taaaattaaa	2100
gtcagtatga	actataataa	atatctgtag	cttaaaaagt	agggtcacat	gtacaggtaa	2160
gtatatgtgt	tattttctgt	cattttctgt	tcataagagt	gtataataaa	acatgattgc	2220

ttaaaaaactt gtatagtgtg ctagatttct gcacctgaat gtatgtttga tgcttttgatt 2280
 tgaaaatggt cttccctgtt atttacattc cgggtgggttt ttaaaattct tacctccatc 2340
 atgcaatttt gaaaattgtg tccagaatta aaagtgcata gaaatagcct ttacaattgt 2400
 agcatggacc tttaaaaatt gttttaaaat cttattttaa. tttaaaccag aagctgaaaa 2460
 atagatcagc tttattatac acaaaattat tactgcttat ctttgctctt ttccttggtta 2520
 tcccgcgaagg tttagttgag aagatacaaa atgttttacag tgttggcact tagagttttt 2580
 aaattcaagt acatgaaatt cagtaatagc attgccttga gctaactagg aagtaccggg 2640
 aaaaaagtta aatctacatc aagtttcttt tgaactttga agtgttttct gaccactgc 2700
 taactgtagc aacaaaattt 2720

<210> 13
 <211> 2698
 <212> DNA
 <213> Homo sapiens

<400> 13
 gagctgaaat tcgcgggtgcg acgggaggga gtggagaagg aggtgagggg gccaggatc 60
 ggggggcgcc ctgaggcaag gggacgccgg tgggtcgaag cgcagcccg cggccgcagg 120
 ctcggctccg cactgcgcg cctcccggtc tcctcgctc gggcgccgag gcaggagag 180
 aatgagcccc gggacccgcc gggggacggc ccgggcccagg cccgggatct agaacggccg 240
 tagggggaag ggagccgcc tccccacggc gccttttcgg aactgccgtg gactcgagga 300
 cgctgggtgc cggcctccta gggctgtgct gttttgtttt gaccctcgca ttgtgcagaa 360
 ttaaagtgca gtaaaatgtc cactaggacc ccattgccaa cgggtgaatga acgagacact 420
 gaaaaccaca cgtcacatgg agatgggct caagaagtta cctctcgta cagccgctca 480
 ggagctcggg ttagaaactc tatagcctcc tgtgcagatg aacaacctca catcggaac 540
 tacagactgt tgaaaacaat cggcaagggg aattttgcaa aagtaaaatt ggcaagacat 600
 atccttacag gcagagaggt tgcaataaaa ataattgaca aaactcagtt gaatccaaca 660
 agtctacaaa agctcttcag agaagtaaga ataatgaaga ttttaaata tcccaatata 720
 gtgaagttat tcgaagtcac tgaaactgaa aaaacactct acctaatacat ggaatatgca 780
 agtggagggtg aagtatttga ctatttggtt gcacatggca ggatgaagga aaaagaagca 840
 agatctaaat ttagacagat tgtgtctgca gttcaatact gccatcagaa acggatcgta 900
 catcgagacc tcaaggctga aaatctattg ttagatgccg atatgaacat taaaatagca 960
 gatttcggtt ttagcaatga atttactgtt ggcggtaaac tcgacacgtt ttgtggcagt 1020
 cctccatagc cagcacctga gctcttcag ggcaagaaat atgacggggc agaagtggat 1080
 gtgtggagtc tgggggtcat ttatacaca ctagtcagtg gctcacttcc ctttgatggg 1140

caaaacctaa aggaactgag agagagagta ttaagaggga aatacagaat tcccttctac 1200
 atgtctacag actgtgaaaa ctttctcaaa cgtttcctgg tgctaaatcc aattaaacgc 1260
 ggcactctag agcaaatcat gaaggacagg tggatcaatg cagggcatga agaagatgaa 1320
 ctcaaaccat ttgttgaacc agagctagac atctcagacc aaaaaagaat agatattatg 1380
 gtgggaatgg gatattcaca agaagaaatt caagaatctc ttagtaagat gaaatacagat 1440
 gaaatcacag ctacatattt gttattgggg agaaaatctt cagagctgga tgctagtgat 1500
 tccagttcta gcagcaatct ttcacttgct aaggttaggc cgagcagtga tctcaacaac 1560
 agtactggcc agtctcctca ccacaaagtg cagagaagtg tttcttcaag ccaaaagcaa 1620
 agacgctaca gtgaccatgc tggaccagct attccttctg ttgtggcgta tccgaaaagg 1680
 agtcagacaa gcactgcaga tggtgacctc aaagaagatg gaatttcctc ccggaaatca 1740
 agtggcagtg ctgttggagg aaaggggaatt gctccagcca gtcccatgct tgggaatgca 1800
 agtaatccta ataaggcgga tattcctgaa cgcaagaaaa gctccactgt ccctagtagt 1860
 aacacagcat ctggtggaat gacacgacga aatacttatg tttgcagtga gagaactaca 1920
 gctgatagac actcagtgat tcagaatggc aaagaaaaca gcactattcc tgatcagaga 1980
 actccagttg cttcaacaca cagtatcagt agtgcagcca cccagatcg aatccgcttc 2040
 ccaagaggca ctgccagtcg tagcactttc caccggccagc cccggggaacg gcgaaccgca 2100
 acatataatg gccctcctgc ctctcccagc ctgtcccatg aagccacacc attgtcccag 2160
 actcgaagcc gaggtccac taatctcttt agtaaattaa cttcaaaact cacaaggagt 2220
 cgcaatgtat ctgctgagca aaaagatgaa aacaaagaag caaagcctcg atccctacgc 2280
 ttcacctgga gcatgaaaac cactagttca atggatcccg gggacatgat gcgggaaatc 2340
 cgcaaagtgt tggacgcaa taactgcgac tatgagcaga gggagcgctt cttgctcttc 2400
 tgcgtccacg gagatgggca cgcgggagaa ctcgtgcagt gggaaatgga agtgtgcaag 2460
 ctgccaagac tgtctctgaa cggggtccgg ttaagcgga tatcggggac atccatagcc 2520
 ttcaaaaata ttgcttccaa aattgccaat gagctaaagc tgtaaccag tgattatgat 2580
 gtaaattaag tagcaagtaa agtgttttcc tgaacactga tggaaatgta tagaataata 2640
 tttaggcaat aacgtctgca tcttctaaat catgaaatta aagtctgagg acgagagc 2698

<210> 14

<211> 2914

<212> DNA

<213> Homo sapiens

<400> 14

gacggccccg gccaggcccc ggatctagaa cggccgtagg ggggaaggag ccgccctccc 60

cacggcgctt tttcggaact gccgtggact cgaggacgct ggtcgccggc ctccctagggc 120
 tgtgctgttt tgttttgacc ctgcattgt gcagaattaa agtgcagtaa aatgtccact 180
 aggaccccat tgccaacggt gaatgaacga gacactgaaa accacacgtc acatggagat 240
 gggcgtcaag aagttacctc tcgtaccagc cgctcaggag ctcggtgtag aaactctata 300
 gcctcctgtg cagatgaaca acctcacatc ggaaactaca gactgttgaa aacaatcggc 360
 aaggggaatt ttgcaaaagt aaaattggca agacatatcc ttacaggcag agagggttgca 420
 ataaaaataa ttgacaaaac tcagttgaat ccaacaagtc tacaaaagct cttcagagaa 480
 gtaagaataa tgaagatttt aaatcatccc aatatagtga agttattcga agtcattgaa 540
 actcaaaaaa cactctacct aatcatggaa tatgcaagtg gaggtaaagt atttgactat 600
 ttggttgacac atggcaggat gaaggaaaaa gaagcaagat ctaaatttag acagattgtg 660
 tctgcagttc aatactgcca tcagaaacgg atcgtaacac gagacctcaa ggctgaaaat 720
 ctattgttag atgccgatat gaacattaaa atagcagatt tcgggttttag caatgaattt 780
 actgttggcg gtaaaactoga cacgttttgt ggcagtcctc catacgagc acctgagctc 840
 ttccagggca agaaatatga cgggccagaa gtggatgtgt ggagtctggg ggtcatttta 900
 tacacactag tcagtggctc acttcccttt gatgggcaaa acctaaagga actgagagag 960
 agagtattaa gagggaaata cagaattccc ttctacatgt ctacagactg tgaaaacctt 1020
 ctcaaacggt tcctggtgct aaatccaatt aaacgcggca ctctagagca aatcatgaag 1080
 gacaggtgga tcaatgcagg gcatgaagaa gatgaactca aaccatttgt tgaaccagag 1140
 ctagacatct cagacccaaa agaatagat attatggtgg gaatgggata ttcacaagaa 1200
 gaaattcaag aatctcttag taagatgaaa tacgatgaaa tcacagctac atatttgtta 1260
 ttggggagaa aatcttcaga ggtagggccg agcagtgatc tcaacaacag tactggccag 1320
 tctcctcacc acaaagtgcg gagaagtgtt tcttcaagcc aaaagcaaag acgctacagt 1380
 gaccatgctg gaccaggtat tccttctgtt gtggcgatc cgaaaaggag tcagaccagc 1440
 actgcagata gtgacctcaa agaagatgga atttcctccc ggaaatcaac tggcagtgct 1500
 gttggaggaa agggaattgc tccagccagt cccatgcttg ggaatgcaag taatccta 1560
 aaggcggata ttcctgaacg caagaaaagc tccactgtcc ctagtagtaa cacagcatct 1620
 ggtggaatga cacgacgaaa tacttatgtt tgcagtgaga gaactacaga tgatagacac 1680
 tcagtgattc agaattggca agaaaacagc actattcctg atcagagaa cccagttgct 1740
 tcaacacaca gtatcagtag tgcagccacc ccagatcgaa tccgcttccc aagaggca 1800
 gccagtcgta gcactttcca cggccagccc cgggaacggc gaaccgcaac atataatggc 1860
 cctcctgcct ctcccagcct gtcccatgaa gccacacat tgtcccagac tcgaagccga 1920

```

ggctccacta ctctcttttag taaattaact tcaaaaactca caaggagtcg caatgtatct 1980
gctaagcaaa aagatgaaaa caaagaagca aagcctcgat ccctacgctt cacctggagc 2040
atgaaaacca ctagttcaat ggatcccggg gacatgatgc gggaaatccg caaagtgttg 2100
gacgccataa actgcgacta tgagcagagg gagcgcttct tgctcttctg cgtccacgga 2160
gatgggcacg cggagaacct cgtgcagtgg gaaatggaag tgtgcaagct gccaaagactg 2220
tctctgaacg ggggtccggtt taagcggata tcggggacat ccatagcctt caaaaatatt 2280
gcttccaaaa ttgccaatga gctaaagctg taaccacgtg attatgatgt aaattaagta 2340
gcaagtaaag tgttttctctg aacactgatg gaaatgtata gaataatatt taggcaataa 2400
cgtctgcata ttctaaatca tgaaattaaa gtctgaggac gagagcacgc ctggggagcga 2460
aagctggcct tttttctacg aatgcactac attaaagatg tgcaacctat gcgccccctg 2520
ccctacttcc gttaccctga gagtcggcgt gtggcccat ctccatgtgc ctcccgctg 2580
gggtgggtgtg agagtggacg gtatgtgtgt gaagtgggtg atatggaagc atctccctac 2640
actggcagcc agtcattact agtacctctg cgggagatca tccggtgcta aaacattaca 2700
gttgccaagg aggaaaatac tgaatgactg ctaagaatta accttaagac cagttcatag 2760
ttaatacagg tttaacagttc atgcctgtgg tttgtgtttt gttgttttgt gtttttttag 2820
tgcaaaaggt ttaaatattat agttgtgaac attgcttggtg tgtgtttttc taagtagatt 2880
cacaagataa ttaaaaattc actttttctc aggt 2914

```

```

<210> 15
<211> 3895
<212> DNA
<213> Homo sapiens

```

```

<400> 15
ctgcaggaat tccgatcctt ccgcagggtc acctacggaa accttggtac gacttttact 60
tctctagat agtcaagttc gaccgtcttc tcagcgctcc gccagggccg tggggccgacc 120
ccggcggggc cgatccgagg gcctcactaa accatccaat cggtagtagc gacgggcggt 180
gtgtacaaag ggcagggact taatcaacgc aagcttatga cccgcactta ctgggaattc 240
ctcgttcatg gggaataatt gcaatccccg atccccatca cgaatggggg tcaacggggt 300
acccgcgcct gccggcgtag ggtaggcaca cgctgagcca gtcagtgtag cgcgcgtgca 360
gccccggaca tctaagggca tcacagacct gtatttgctc aatctcgggt ggctgaacgc 420
cacttgctcc tctaagaagt tgggggacgc cgaccgctcg ggggtcgcgt aactagttag 480
catgccagag tctcgttcgt tatcggaatt aaccagacaa atcgctccac caactaagaa 540
cggccatgca ccaccaccca cggaatcgag aaagagctat caatctgtca atcctgtccg 600
tgtccggggc ggggtgaggtt tcccggtgtg agtcaaatta agccgcaggc tccactcctg 660

```

gtggtgccct tccgtcaatt cctttaagtt tcagctttgc aaccatactc cccccggaac	720
ccaaagactt tggtttcccg gaagctgccc ggcgggtcat gggaataacg ccgccgcac	780
gccggtcggc atcgtttatg gtcggaacta cgacggtatc tgatcgctctt cgaacctccg	840
actttcgttc ttgattaatg aaaacattct tggcaaagtc tttcgctctg gtcgctcttg	900
cgccggtcca agaatttcgg aattccgcag cggcggccag cagggcgagg gctgaggcag	960
caagctcgct agagaggag aagcagtcgg gcgcaggcgc ctctccgca gccgctcca	1020
tggtcggcgc ccacagcccg cggcggcctg tcttgcgctc cacttccttc acatcctcct	1080
ccgcctcctc gttttcaggc gccgcggcg gcgctgtgtg gaggcccgcg agctgaaatt	1140
cgcggtgcca cgggaggag tggagaagga ggtgagggg cccaggatcg cggggcgccc	1200
tgaggcaagg ggacgccggc gggccgaagc gcagcccgcc gcccgaggc tcggctccgc	1260
cactgccgcc ctcccggtct cctgcctcg gccgccagg cagggagaga atgagccccg	1320
ggaccccgcg ggggacggc cgggccaggc ccgggatcta gacggcgta gggggaaggg	1380
agccgcctc cccacggcg cttttcgga ctgccgtgga ctcgaggacg ctggtcgccg	1440
gcctcctagg gctgtgctgt tttgttttga cctcgcatt gtgcagaatt aaagtgcagt	1500
aaaatgtcca ctaggacccc attgccaacg gtgaatgaac gagacactga aaaccacacg	1560
tcacatggag atgggcgtca agaagttacc tctcgtaacca gccgctcagg agctcgggtg	1620
agaaactcta tagcctcctg tgcagatgaa caacctcaca tcggaaacta cagactgttg	1680
aaaacaatcg gcaaggggaa ttttgcaaaa gtaaaattgg caagacatat ccttacaggc	1740
agagagggtg caataaaaat aattgacaaa actcagttga atccaacaag tctacaaaag	1800
ctcttcagag aagtaagaat aatgaagatt ttaaatactc ccaatatagt gaagttattc	1860
gaagtcattg aaactgaaaa aacactctac ctaatcatgg aatatgcaag tggagggtgaa	1920
gtatttgact atttggttgc acatggcaag atgaaggaaa aagaagcaag atctaaattt	1980
agacagggtt gtcaagctgg acagactatt aaagttcaag tctcctttga tttgcttagt	2040
ctgatgttta cattttattgt gtctgcagtt caatactgcc atcagaaacg gatcgtaac	2100
cgagacctca aggctgaaaa tctattgtta gatgccgata tgaacattaa aatagcagat	2160
ttcggtttta gcaatgaatt tactgttggc ggtaaactcg acacgttttg tggcagtcct	2220
ccatacgag cacctgagct cttccagggc aagaaatatg acgggccaga agtggatgtg	2280
tggagtctgg gggtcatttt atacacacta gtcagtggt cacttccttt tgatgggcaa	2340
aacctaaagg aactgagaga gagagtatta agagggaat acagaattcc cttctacatg	2400
tctacagact gtgaaaaact tctcaaactg ttcctggtgc taaatccaat taaacgcggc	2460
actctagagc aaatcatgaa ggacagggtg atcaatgcag ggcataaga agatgaactc	2520

```

aaaccatttg ttgaaccaga gctagacatc tcagacccaa aaagaataga tattatgggtg 2580
ggaatgggat attcacaaga agaaattcaa gaatctctta gtaagatgaa atacgatgaa 2640
atcacagcta catatttggtt attggggaga aaatcttcag agctggatgc tagtgattcc 2700
agttctagca gcaatctttc acttgctaag gttaggccga gcagtgatct caacaacagt 2760
actggccagt ctctcacca caaagtgcag agaagtgttt cttcaagcca aaagcaaaga 2820
cgctacagtg accatgctgg accagctatt ccttctgttg tggcgtatcc gaaaaggagt 2880
cagacaagca ctgcagatgg tgacctcaa gaagatggaa tttcctccc gaaatcaagt 2940
ggcagtgtg ttggaggaaa gggaattgct ccagccagtc ccatgcttgg gaatgcaagt 3000
aatcctaata agggggatat tctgaacgc aagaaaagct cactgtccc tagtagtaac 3060
acagcatctg gtggaatgac acgacgaaat acttatgttt gcagtgagag aactacagct 3120
gatagacact cagtgattca gaatggcaaa gaaaacagca ctattcctga tcagagaact 3180
ccagttgctt caacacacag tatcagtagt gcagccaccc cagatcgaat ccgcttccca 3240
agaggcactg ccagtcgtag cactttccac ggccagcccc gggaacggcg aaccgcaaca 3300
tataatggcc ctctgcctc tcccagcctg tcccatgaag ccacaccatt gtcccagact 3360
cgaagccgag gctccactaa tctcttttagt aaattaactt caaaactcac aaggagtgcg 3420
aatgtatctg ctgagcaaaa agatgaaaac aaagaagcaa agcctcgatc cctacgcttc 3480
acctggagca tgaaaaccac tagttcaatg gatccccggg acatgatgcg ggaaatccgc 3540
aaagtgttgg acgccaataa ctgcgactat gagcagaggg agcgcttctt gctcttctgc 3600
gtccacggag atgggcacgc ggagaacctc gtgcagtggg aaatggaagt gtgcaagctg 3660
ccaagactgt ctctgaacgg ggtccggttt aagcggatat cggggacatc catagccttc 3720
aaaaatattg cttccaaaat tgccaatgag ctaaagctgt aaccagtgta ttatgatgta 3780
aattaagtag caagtaaagt gttttcctga aactgatgg aaatgtatag aataatattt 3840
aggcaataac gtctgcatct tctaaatcat gaaattaaag tctgaggacg agagc 3895

```

<210> 16

<211> 2145

<212> DNA

<213> Homo sapiens

<400> 16

```

atgtccacta ggacccatt gccaacggtg aatgaacgag aactgaaaa ccacacgtca 60
catggagatg ggcgtcaaga agttacctct cgtaccagcc gctcaggagc tcggtgtaga 120
aactctatag ctcctgtgc agatgaacaa cctcacatcg gaaactacag actgttgaaa 180
acaatcggca aggggaattt tgcaaaagta aaattggcaa gacatatcct tacaggcaga 240

```

gagggttgcaa taaaaataat tgacaaaact cagttgaatc caacaagtct acaaaagctc	300
ttcagagaag taagaataat gaagatttta aatcatocca atatagttaa gttattcgaa	360
gtcattgaaa ctgaaaaaac actctaccta atcatggaat atgcaagtgg aggtgaagta	420
tttgactatt tgggtgcaca tggcaggatg aaggaaaaag aagcaagatc taaatttaga	480
cagattgtgt ctgcagttca atactgccat cagaaacgga tcgtacatcg agacctcaag	540
gctgaaaatc tattgttaga tgccgatatg aacattaaaa tagcagattt cggtttttagc	600
aatgaattta ctgttggcgg taaactcgac acgttttgtg gcagtcctcc atacgcagca	660
cctgagctct tccagggcaa gaaatatgac gggccagaag tggatgtgtg gagtctgggg	720
gtcatttttat acacactagt cagtggctca cttccctttg atgggcaaaa cctaaaggaa	780
ctgagagaga gagtattaag agggaaatac agaattccct tctacatgtc tacagactgt	840
gaaaaccttc tcaaacgttt cctgggtgcta aatccaatta aacgcggcac tctagagcaa	900
atcatgaagg acaggtggat caatgcaggg catgaagaag atgaactcaa accatttgtt	960
gaaccagagc tagacatctc agaccaaaaa agaatagata ttatgggtggg aatgggatat	1020
tcacaagaag aaattcaaga atctcttagt aagatgaaat acgatgaaat cacagctaca	1080
tattttgttat tggggagaaa atcttcagag gttaggccga gcagtgatct caacaacagt	1140
actggccagt ctctcacca caaagtgcag agaagtgttt cttcaagcca aaagcaaaga	1200
cgctacagtg accatgctgg accagctatt ccttctgttg tggcgatatcc gaaaaggagt	1260
cagaccagca ctgcagatag tgacctcaaa gaagatggaa tttcctcccg gaaatcaagt	1320
ggcagtgtgt ttggaggaaa gggaattgct ccagccagtc ccatgcttgg gaatgcaagt	1380
aatcctaata aggcggatat tcctgaacgc aagaaaagct ccactgtccc tagtagtaac	1440
acagcatctg gtggaatgac acgacgaaat acttatgttt gcagtgagag aactacagct	1500
gatagacact cagtgattca gaatggcaaa gaaaacagca ctattcctga tcagagaact	1560
ccagttgctt caacacacag tatcagtagt gcagccaccc cagatogaat ccgcttccca	1620
agaggcactg ccagtcgtag cactttccac ggccagcccc gggaacggcg aaccgcaaca	1680
tataatggcc ctctgcctc tcccagcctg tcccatgaag ccacaccatt gtcccagact	1740
cgaagccgag gctccactaa tctcttttagt aaattaactt caaaactcac aaggagtgc	1800
aatgtatctg ctgagcaaaa agatgaaaac aaagaagcaa agcctcgatc cctacgcttc	1860
acctggagca tgaaaaccac tagttcaatg gatccggggg acatgatgag ggaaatccgc	1920
aaagtgttgg acgccaataa ctgcgactat gagcagaggg agcgcttctt gctcttctgc	1980
gtccacggag atgggcacgc ggagaacctc gtgcagtggg aaatggaagt gtgcaagctg	2040
ccaagactgt ctctgaacgg ggtccggttt aagcggatat cggggacatc catagccttc	2100

aaaaatattg cttccaaaat tgccaatgag ctaaagctgt aacc 2145

<210> 17

<211> 2193

<212> DNA

<213> Homo sapiens

<400> 17

atgtccacta ggacccatt gccaacggtg aatgaacgag aactgaaaa ccacacgtca 60
catggagatg ggcgtcaaga agttacctct cgtaccagcc gctcaggagc tcggtgtaga 120
aactctatag cctcctgtgc agatgaacaa cctcacatcg gaaactacag actggttga 180
acaatoggca aggggaattt tgcaaaagta aaattggcaa gacatatcct tacaggcaga 240
gaggttgcaa taaaaataat tgacaaaact cagttgaatc caacaagtct acaaaagctc 300
ttcagagaag taagaataat gaagatttta aatcatccca atatagtga gttattcgaa 360
gtcattgaaa ctgaaaaaac actctacctt atcatggaat atgcaagtgg aggtgaagta 420
tttgactatt tgggtgcaca tggcaggatg aaggaaaaag aagcaagatc taaatttaga 480
cagatttgtt ctgcagttca atactgccat cagaaacgga tcgtacatcg agacctcaag 540
gctgaaaatc tattgttaga tgccgatatg aacattaaaa tagcagattt cggtttttagc 600
aatgaattta ctgttggcgg taaactcgac acgtttttgtg gcagtcctcc atacgcagca 660
cctgagctct tccagggcaa gaaatatgac gggccagaag tggatgtgtg gaggctgggg 720
gtcattttat acacactagt cagtggctca cttccctttg atgggcaaaa cctaaaggaa 780
ctgagagaga gagtattaag agggaaatac agaattccct tctacatgtc tacagactgt 840
gaaaaccttc tcaaacgttt cctggtgcta aatccaatta aacgcggcac tctagagcaa 900
atcatgaagg acaggtggat caatgcaggg catgaagaag atgaactcaa accatttgtt 960
gaaccagagc tagacatctc agaccaaaaa agaatagata ttatgggtggg aatgggatat 1020
tcacaagaag aaattcaaga atctcttagt aagatgaaat acgatgaaat cacagctaca 1080
tatttgttat tggggagaaa atcttcagag ctggatgcta gtgattccag ttctagcagc 1140
aatctttcac ttgctaaggt taggccgagc agtgatctca acaacagtac tggccagtct 1200
cctcaccaca aagtgcagag aagtgtttct tcaagccaaa agcaaagacg ctacagtgc 1260
catgctggac cagctattcc ttctgttgtg gcgtatccga aaaggagtca gaccagcact 1320
gcagatagtg acctcaaaga agatggaatt tcctcccga aatcaagtgg cagtgcgtgtt 1380
ggaggaaagg gaattgctcc agccagtccc atgcttggga atgcaagtaa tctaataag 1440
gcggatattc ctgaacgcaa gaaaagctcc actgtcccta gtagtaaacac agcatctggt 1500
ggaatgacac gacgaaatac ttatgtttgc agtgagagaa ctacagctga tagacactca 1560
gtgattcaga atggcaaaga aaacagcact attcctgatc agagaactcc agttgcttca 1620

acacacagta tcagtagtgc agccacccca gatcgaatcc gcttcccaag aggcactgcc 1680
 agtcgtagca ctttccacgg ccagccccgg gaacggcgaa ccgcaacata taatggccct 1740
 cctgcctctc ccagcctgtc ccatgaagcc acaccattgt cccagactcg aagccgaggg 1800
 tccactaatc tctttagtaa attaaacttca aaactcaciaa ggagtcgcaa tgtatctgct 1860
 gagcaaaaag atgaaaacaa agaagcaaag cctcgatccc tacgcttcac ctggagcatg 1920
 aaaaccacta gttcaatgga tcccgggggac atgatgcggg aaatccgcaa agtggttgac 1980
 gccataact gcgactatga gcagagggag cgcttcttgc tcttctgcgt ccacggagat 2040
 gggcacgcgg agaacctcgt gcagtgggaa atggaagtgt gcaagctgcc aagactgtct 2100
 ctgaacgggg tccggtttaa gcggatatcg gggacatcca tagccttcaa aaatattgct 2160
 tccaaaattg ccaatgagct aaagctgtaa ccc 2193

<210> 18

<211> 3373

<212> DNA

<213> Homo sapiens

<400> 18

caggcgctc ctccgcagcc cgctccatgg tcggcgccca cagcccgcg cggcctgtct 60
 tgcgctccac ttccctcaca tctcctccg cctcctcgtt ttcaggcgcc gccggcgcg 120
 ctgtgtggag gcccgcgagc tgaaattcgc ggtgcgacgg gagggagtgg agaaggaggt 180
 gagggggccc aggatcgcg ggcgccctga ggcaagggga cgccggcggg ccgaagcgca 240
 gcccgccgcc cgcaggetcg gctccgccac tgccgcctc ccggtctcct cgcctcggcc 300
 gccgaggcag ggagagaatg agccccggga cccgccgggg acggcccggg ccaggcccg 360
 gatctagaac ggccgtaggg ggaaggagc cgccctcccc acggcgccct ttcggaactg 420
 ccgtggactc gaggacgctg gtcgccggcc tcctagggct gtgctgtttt gttttgacct 480
 tcgcattgtg cagaattaaa gtgcagtaaa atgtccacta ggacccatt gccaacgggtg 540
 aatgaacgag aactgaaaa ccacacgtca catggagatg ggcgtcaaga agttacctct 600
 cgtaccagcc gctcaggagc tcggtgtaga aactctatag cctcctgtgc agatgaacaa 660
 cctcacatcg gaaactacag actgttgaaa acaatcgga aggggaattt tgcaaaaagta 720
 aaattggcaa gacatatcct tacaggcaga gaggttgcaa taaaaataat tgacaaaact 780
 cagttgaatc caacaagtct acaaaagctc ttcagagaag taagaataat gaagatttta 840
 aatcatccca atatagtga gttattcgaa gtcattgaaa ctgaaaaaac actctaccta 900
 atcatggaat atgcaagtgg aggtaaagta tttgactatt tggttgca tggcaggatg 960
 aaggaaaaag aagcaagatc taaatttaga cagattgtgt ctgcagttca atactgcat 1020

cagaaacgga tcgtacatcg agacctcaag gctgaaaatc tattgttaga tgccgatatg 1080
 aacattaaaa tagcagattt cggtttttagc aatgaattta ctgttggcgg taaactcgac 1140
 acgttttgtg gcagtcctcc atacgcagca cctgagctct tccagggcaa gaaatatgac 1200
 gggccagaag tggatgtgtg gagtctgggg gtcattttat acacactagt cagtggctca 1260
 cttccctttg atgggcaaaa cctaaaggaa ctgagagaga gagtattaag agggaaatac 1320
 agaattccct tctacatgtc tacagactgt gaaaaccttc tcaaacgttt cctgggtgcta 1380
 aatccaatta aacgcggcac tctagagcaa atcatgaagg acaggtggat caatgcaggg 1440
 catgaagaag atgaactcaa accatttgtt gaaccagagc tagacatctc agacaaaaaa 1500
 agaatagata ttatgggtggg aatgggatat tcacaagaag aaattcaaga atctcttagt 1560
 aagatgaaat acgatgaaat cacagctaca tatttgttat tggggagaaa atcttcagag 1620
 ctggatgcta gtgattccag ttctagcagc aatctttcac ttgctaaggt taggccgagc 1680
 agtgatctca acaacagtac tggccagtct cctcaccaca aagtgcagag aagtgtttct 1740
 tcaagccaaa agcaaagacg ctacagtgc catgctggac cagctattcc ttctgttgtg 1800
 gcgtatccga aaaggagtca gaccagcact gcagatagtg acctcaaaga agatggaatt 1860
 tcctcccggg aatcaagtgg cagtgcgtgtt ggaggaaagg gaattgctcc agccagtccc 1920
 atgcttggga atgcaagtaa tcctaataag gcggatattc ctgaacgcaa gaaaagctcc 1980
 actgtcccta gtagtaacac agcatctggt ggaatgacac gacgaaatac ttatgtttgc 2040
 agtgagagaa ctacagctga tagacactca gtgattcaga atggcaaaga aaacagcact 2100
 attcctgatc agagaactcc agttgcttca acacacagta tcagtagtgc agccacccca 2160
 gatcgaatcc gcttcccaag aggcaactgcc agtcgtagca ctttccacgg ccagccccgg 2220
 gaacggcgaa ccgcaacata taatggccct cctgcctctc ccagcctgtc ccatgaagcc 2280
 acaccattgt ccagactcg aagccgaggg tcactaatac tcttttagtaa attaaattca 2340
 aaactcacia ggagaaacat gtcattcagg tttatcaaaa ggcttccaac tgaatatgag 2400
 aggaacggga gatatgaggg ctcaagtgcg aatgtatctg ctgagcaaaa agatgaaaac 2460
 aaagaagcaa agcctcgatc cctacgcttc acctggagca tgaaaaccac tagttcaatg 2520
 gatccccggg acatgatgcg ggaaatccgc aaagtgttgg acgccaataa ctgcgactat 2580
 gagcagaggg agcgcttctt gctcttctgc gtccacggag atgggcacgc ggagaacctc 2640
 gtgcagtggg aaatggaagt gtgcaagctg ccaagactgt ctctgaacgg ggtccggttt 2700
 aagcggatat cggggacatc catagccttc aaaaatattg cttccaaaat tgccaatgag 2760
 ctaaagctgt aaccagtgta ttatgatgta aattaagtag caagtaaagt gttttcctga 2820
 acactgatgg aaatgtatag aataatattt aggcaataac gtctgcatct tctaaatcat 2880

```

gaaattaaag tctgaggacg agagcacgcc tgggagcgaa agctggcctt ttttctacga 2940
atgcactaca ttaaagatgt gcaacctatg cgccccctgc cctacttccg ttaccctgag 3000
agtcggcgtg tggccccatc tccatgtgcc tcccgctctgg gtgggtgtga gagtggacgg 3060
tatgtgtgtg aagtgggtga tatggaagca tctccctaca ctggcagcca gtcattacta 3120
gtacctctgc gggagatcat ccggtgctaa aacattacag ttgccaagga ggaaaatact 3180
gaatgactgc taagaattaa ccttaagacc agttcatagt taatacaggt ttacagttca 3240
tgcctgtggg tttgtgtttg ttgttttgtg tttttttagt gcaaaagggt taaatttata 3300
gttgtgaaca ttgcttgtgt gtgtttttct aagtagattc acaagataat taaaaattca 3360
ctttttctca ggt 3373

```

```

<210> 19
<211> 3609
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (3606)..(3606)
<223> "n" is A, C, G, or T

```

```

<400> 19
cgctccctc cgccgcgcgt tgggcccggc ccgcgcctcc tccgcggccc ccgcccgcct 60
gcctgcccgc cgcccccatg ggcgccgggg tcccgcgtgc acggggccac taggaccctc 120
ggcgccctt cccctccccc gccctgcccc ctctcccgcc gcgcgggacc gggcgttctc 180
ggcgcccagc ttttgagctc gcgtccccag gccggcgggg ggggagggga agagagggga 240
ccctgggacc ccgccccccc ccacccggcc gccctgccc ccggggacc ggagaagatg 300
tcttcgcgga cgggtgctgg cccgggcaac gatcggaact cggacacgca tggcaccttg 360
ggcagtggcc gctcctcgga caaaggcccg tcctgggtcca gccgctcact gggtgcccgt 420
tgccggaact ccatgcctc ctgtcccag gagcagcccc acgtgggcaa ctaccgcctg 480
ctgaggacca ttgggaaggg caactctgcc aaagtcaagc tggctcggca catcctcact 540
ggtcgggaggt ttgccatcaa gattatcgac aaaaccagc tgaatcccag cagcctgcag 600
aagctgttcc gagaagtccg catcatgaag ggcctaaacc accccaacat cgtgaagctc 660
tttgaggtga ttgagactga gaagacgctg tacctgggtga tggagtacgc aagtgtctga 720
gaagtgtttg actacctcgt gtcgcatggc cgcgtgaagg agaaggaagc tcgagccaag 780
ttccgacaga ttgtttcggc tgtgcactat tgtcaccaga aaaatatgt acacagggac 840
ctgaaggctg agaacctctt gctggatgcc gaggccaaca tcaagattgc tgactttggc 900
ttcagcaacg agttcacgct gggatcgaa ctggacacgt tctgcgggag cccccatat 960

```

gccgccccgg agctgtttca gggcaagaag tacgacgggc cggaggtgga catctggagc 1020
 ctggggagtca tctgtacac cctcgtcagc ggctccctgc ccttcgacgg gcacaacctc 1080
 aaggagctgc gggagcgagt actcagaggg aagtaccggg tccctttcta catgtcaaca 1140
 gactgtgaga gcatcctgcg gagatTTTTg gtgctgaacc cagctaaacg ctgtactctc 1200
 gagcaaatca tgaaagacaa atggatcaac atcggctatg aggggtgagga gttgaagcca 1260
 tacacagagc ccgaggagga cttcggggac accaagagaa ttgaggtgat ggtgggtatg 1320
 ggctacacac gggaagaaat caaagagtcc ttgaccagcc agaagtacaa cgaagtgacc 1380
 gccacctacc tctgtctggg caggaagact gaggaggggtg gggaccgggg cggcccaggg 1440
 ctggccctgg cacgggtgcg ggcgcccagc gacaccacca acggaacaag ttccagcaaa 1500
 ggcaccagcc acagcaaagg gcagcggagt tctcttcca cctaccaccg ccagcgcagg 1560
 catagcgatt tctgtggccc atccccctgca cccctgcacc ccaaaccgag cccgacgagc 1620
 acggggggagg cggagctgaa ggaggagcgg ctgccaggcc ggaaggcgag ctgcagcacc 1680
 gcgggggagtg ggagtcgagg gctgcccccc tccagcccca tggtcagcag cggccacaac 1740
 cccaacaagg cagagatccc agagcggcgg aaggacagca cgagcaccac caacaacctc 1800
 cctcctagca tgatgacctg cagaaacacc tacgtttgca cagaacgccc gggggctgag 1860
 cgcccgctac tgttgccaaa tgggaaagaa aacagctcag gcacccacag ggtgccccct 1920
 gcctccccct ccagtcacag cctggcacc ccatcagggg agcggagccg cctggcacgc 1980
 ggttcacca tccgcagcac cttccatggt ggccaggtcc gggaccggcg ggcagggggt 2040
 gggggtggtg ggggtgtgca gaatgggccc cctgcctctc ccacactggc ccatgaggct 2100
 gcacccctgc ccgccggcg gccccgccc accaccaacc tcttcaccaa gctgacctcc 2160
 aaactgacct gaagggttac cctcgatccc tctaaacggc agaactctaa ccgctgtgtt 2220
 tcgggcgcct ctctgcccc aaggatccaag atcaggtcgc agacgaacct gagagaatcg 2280
 ggggacctga ggtcacaagt tgccatctac cttgggatca aacggaaacc gcccccggc 2340
 tgctccgatt cccctggagt gtgaagctga ccagctcgcg ccctcctgag gccctgatgg 2400
 cagctctgcg ccaggccaca gcagccgccc gctgcgctg ccgccagcca cagccgttcc 2460
 tgctggcctg cctgcacggg ggtgcggggc ggcccgagcc cctgtccac ttcgaagtgg 2520
 aggtctgcca gctgccccgg ccaggcttgc ggggagttct cttccgccgt gtggcgggca 2580
 ccgccctggc cttccgcacc ctctcacc gcattctcaa cgacctcgag ctctgagcca 2640
 ccacggctcc agggccctta ctcttctct cccttgctgc cttcacttct acaggagggg 2700
 aaggggcccag ggaggggatt ctccctttat catcacctca gtttcctga attatatttg 2760
 ggggcaaaga ttgtccctc tgctgttctc tggggccgct cagcacagaa gaaggatgag 2820

ggggctcagc ggggggagct ggcaccttcc tggagcctcc agccagtcct gtccctccctc 2880
 gccctaccaa gagggcacct gaggagactt tggggacagg gcagggggcag ggaggggaaac 2940
 tgaggaaatc ttccattcct cccaacagct caaaattagg ccttggggcag gggcagggag 3000
 agctgctgag cctaaagact ggagaatctg ggggactggg agtggggggtc agagaggcag 3060
 attccttccc ctcccgctcc ctacgctca aacccccact tcctgccccca ggctggcgcg 3120
 gggcactttg tacaaatcct tgtaaatacc ccacaccctc ccctctgcaa aggtctcttg 3180
 aggagctgcc gctgtcacct acgggttttta agttattaca ccccgaccct cctcctgtca 3240
 gccccctcac ctgcagcctg ttgcccaata aatttaagag agtccccccc tccccaatgc 3300
 tgaccctagg attttccctc cctgccctca cctgcaaag agttaaagaa gaggcgtggg 3360
 aatccaggca gtgggttttct ctttcggagc ctcggttttc tcatctgcag aatgggagcg 3420
 gtgggggtgg gaaggtaagg atggtcgtgg aagaaggcag gatgggaactc ggcctcatcc 3480
 ccgaggcccc agttcctata tggggcccc cattcatcca ctcacactcc cagccaccat 3540
 gttacactgg actctaagcc acttcttact ccagtagtaa atttattgca ataaacaatc 3600
 attganccc 3609

<210> 20
 <211> 2085
 <212> DNA
 <213> Homo sapiens

<400> 20
 agatgtcttc gcggacggtg ctggccccgg gcaacgatcg gaactcggac acgcatggca 60
 ccttggggcag tggccgctcc tcggacaaaag gcccgctctg gtccagccgc tcaactgggtg 120
 cccgttgccg gaactccatc gcctcctgtc ccgaggagca gcccacgtg ggcaactacc 180
 gcctgctgag gaccattggg aagggaact ttgccaaagt caagctggct cggcacatcc 240
 tcaactggtcg ggaggttgcc atcaagatta tcgacaaaac ccagctgaat cccagcagcc 300
 tgcagaagct gttccgagaa gtccgcatca tgaagggcct aaaccacccc aacatcgtga 360
 agctctttga ggtgattgag actgagaaga cgctgtacct ggtgatggag tacgcaagtg 420
 ctggagaagt gtttgactac ctcggtgtgc atggccgcat gaaggagaag gaagctcgag 480
 ccaagttccg acagattgtt tcggctgtgc actattgtca ccagaaaaat attgtacaca 540
 gggacctgaa ggctgagaac ctcttgctgg atgccgagc caacatcaag attgctgact 600
 ttggcttcag caacgagttc acgctgggat cgaagctgga cagtttctgc gggagcccc 660
 catatgccgc cccggagctg ttccagggca agaagtacga cgggccggag gtggacatct 720
 ggagcctggg agtcatcctg tacaccctcg tcagcggctc cctgcccttc gacgggcaca 780

acctcaagga gctgctgggag cgagtactca gaggggaagta ccgggtccct ttctacatgt 840
 caacagactg tgagagcatc ctgctggagat ttttgggtgct gaacccagct aaacgctgta 900
 ctctcgagca aatcatgaaa gacaaatgga tcaacatcgg ctatgagggg gaggagtga 960
 agccatacac agagccccgag gaggacttcg gggacaccaa gagaattgag gtgatgggtg 1020
 gtatgggcta cacacgggaa gaaatcaaag agtccttgac cagccagaag tacaacgaag 1080
 tgaccgccac ctacctcctg ctgggcagga agactgagga ggggtggggac cggggcgccc 1140
 cagggctggc cctggcacgg gtgctgggag ccagcgacac caccaacgga acaagttcca 1200
 gcaaaggcac cagccacagc aaagggcagc ggagttcctc ttccacctac caccgccagc 1260
 gcaggcatag cgattttctgt ggcccatccc ctgcacctct gcaccccaaa cgcagccga 1320
 cgagcacggg ggagggcggag ctgaaggagg agcggctgcc aggccggaag gcgagctgca 1380
 gcaccgctggg gagtggggagt cgagggctgc cccctccag cccatgggtc agcagcgccc 1440
 acaaccccaa caaggcagag atccagagc ggcggaagga cagcacgagc aaccccaaca 1500
 acctccctcc tagcatgatg acccgagaa acacctacgt ttgcacagaa cgcccggggg 1560
 ctgagcgccc gtcactgttg ccaaattgga aagaaaacag ctgagcacc ccacgggtgc 1620
 cccctgcctc cccctccagt cacagcctgg ccccccatc aggggagcgg agccgcctgg 1680
 cagcggttc caccatccgc agcaccttc atgggtggca ggtccgggac cggcgggcag 1740
 ggggtggggg tgggtggggg gtgcagaatg gggccctgc ctctcccaca ctggcccatg 1800
 aggtgcacc cctgcccgc gggcgggccc gccccaccac caacctctc accaagctga 1860
 cctccaaact gacccgaagg gttaccctcg atccctctaa acggcagaac tctaaccgct 1920
 gtgtttcggg cgctctctg cccagggat ccaagatcag gtcgcagacg aacctgagag 1980
 aatcggggga cctgaggtca caagttgcca tctaccttgg gatcaaacgg aaaccgcccc 2040
 ccggctgctc cgattcccct ggagtgtgaa gctgaccagc tcgcg 2085

<210> 21

<211> 2278

<212> DNA

<213> Homo sapiens

<400> 21

agatgtcttc ggggacgggtg ctggccccgg gcaacgatcg gaactcggac acgcatggca 60
 ccttgggcag tggccgctcc tcggacaaag gccgctcctg gtccagccgc tctactgggtg 120
 cccgttgccg gaactccatc gcctcctgtc ccgaggagca gcccacgtg ggcaactacc 180
 gcctgctgag gaccattggg aagggcaact ttgcaaagt caagctggct cggcacatcc 240
 tctactggtc ggaggttgcc atcaagatta tcgacaaaac ccagctgaat cccagcagcc 300
 tgcagaagct gttccgagaa gtccgcatca tgaagggcct aaaccacccc aacatcgtga 360

agctctttga ggtgattgag actgagaaga cgctgtacct ggtgatggag tacgcaagtg 420
 ctggagaagt gtttgactac ctctgtctgc atggccgcac gaaggagaag gaagctcgag 480
 ccaagttccg acagattgtt tcggctgtgc actattgtca ccagaaaaat attgtacaca 540
 gggacctgaa ggctgagaac ctcttgctgg atgccgaggc caacatcaag attgctgact 600
 ttggcttcag caacgagttc acgctgggat cgaagctgga cacttctgc gggagcccc 660
 catatgccgc cccggagctg tttcagggca agaagtacga cgggccggag gtggacatct 720
 ggagcctggg agtcatcctg tacaccctcg tcagcggctc cctgcccttc gacgggcaca 780
 acctcaagga gctgcgggag cgagtactca gaggggaagta ccgggtccct ttctacatgt 840
 caacagactg tgagagcatc ctgcggagat ttttggtgct gaaccagct aaacgctgta 900
 ctctcgagca aatcatgaaa gacaaatgga tcaacatcgg ctatgagggg gaggagttga 960
 agccatacac agagcccag gaggacttcg gggacaccaa gagaattgag gtgatgggtg 1020
 gtatgggcta cacacgggaa gaaatcaaag agtccttgac cagccagaag tacaacgaag 1080
 tgaccgccac ctacctcctg ctgggcagga agactgagga ggggtggggac cggggcgccc 1140
 cagggtctgg cctggcacgg gtgcgggcgc ccagcgacac caccaacgga acaagttcca 1200
 gcaaaggcac cagccacagc aaagggcagc ggagttcctc ttccacctac caccgccagc 1260
 gcaggcatag cgattttctg gggccatccc ctgcacccct gcaccccaaa cgcagcccga 1320
 cgagcacggg ggagggcgag ctgaaggagg agcggctgcc aggccggaag gcgagctgca 1380
 gcaccgcggg gagtgggagt cgagggctgc cccctccag ccccatggtc agcagcgccc 1440
 acaaccccaa caaggcagag atcccagagc ggcggaagga cagcacgagc accccaaca 1500
 acctccctcc tagcatgatg acccgagaa acacctacgt ttgcacagaa cgccggggg 1560
 ctgagcgccc gtcactgttg ccaaattggga aagaaaacag ctcaggcacc ccacgggtgc 1620
 cccctgcctc cccctccagt cacagcctgg caccacctc aggggagcgg agccgcctgg 1680
 caccgggttc caccatccgc agcaccttc atggtggcca ggtccgggac cggcgggcag 1740
 ggggtggggg tgggtggggg gtgcagaatg gggccctgc ctctccaca ctggcccatg 1800
 aggtgcacc cctgcccgc gggcgggccc gcccaccac caacctcttc accaagctga 1860
 cctccaaact gacccgaagg gtcgcagacg aacctgagag aatcggggga cctgaggtca 1920
 caagttgcca tctaccttg gatcaaacgg aaaccgccc cgggtgctc cgattccct 1980
 ggagtgtgaa gctgaccagc tcgcgccctc ctgaggccct gatggcagct ctgcgccagg 2040
 ccacagcagc cgcccgctgc cgctgccgc agccacagcc gttcctgctg gcctgcctgc 2100
 acgggggtgc gggcgggccc gagccctgt cccacttcga agtggagggtc tgccagctgc 2160
 cccggccagg cttgcgggga gttctcttc gccgtgtggc gggcaccgc ctggccttc 2220

gcaccctcgt caccgcgcatc tocaacgacc tcgagctctg agccaccacg gtcccagg 2278

<210> 22

<211> 4917

<212> DNA

<213> Homo sapiens

<400> 22

agaagatgtc ttgcgggacg gtgctggccc cgggcaacga tcggaactcg gacacgcatg 60

gcaccttggg cagtggccgc tcctcggaca aaggcccgtc ctgggtccagc cgctcactgg 120

gtgcccgttg ccggaactcc atcgctctct gtcccgagga gcagccccac gtgggcaact 180

accgcctgct gaggaccatt gggaaggcca actttgccaa agtcaagctg gctcggcaca 240

tcctcactgg tcgggagggt gccatcaaga ttatcgacaa aaccagctg aatcccagca 300

gcctgcagaa gctgttccga gaagtccga tcatgaaggc cctaaaccac cccaacatcg 360

tgaagctctt tgagggtgatt gagactgaga agacgctgta cctgggtgatg gaggtagcaa 420

gtgctggaga agtggttgac tacctcgtgt cgcattggcc catgaaggag aaggaagctc 480

gagccaagtt ccgacagatt gtttcggctg tgcactattg tcaccagaaa aatattgtac 540

acagggacct gaaggctgag aacctcttgc tggatgccga ggccaacatc aagattgctg 600

actttggctt cagcaacgag ttcacgctgg gatcgaagct ggacacgttc tgcgggagcc 660

ccccatatgc cggccgggag ctgtttcagg gcaagaagta cgacggggccg gagggtggaca 720

tctggagcct gggagtcac ctgtacaccc tcgtcagcgg ctccctgccc ttcgacgggc 780

acaacctcaa ggagctgcgg gagcgagtac tcagagggaa gtaccgggtc cctttctaca 840

tgtcaacaga ctgtgagagc atcctgcgga gatTTTTTgtt gctgaaccca gctaaacgct 900

gtactctcga gcaaatcatg aaagacaaat ggatcaacat cggctatgag ggtgaggagt 960

tgaagccata cacagagccc gaggaggact tcggggacac caagagaatt gagggtgatgg 1020

tgggtatggg ctacacacgg gaagaaatca aagagtctt gaccagccag aagtacaacg 1080

aagtgaccgc cactacctc ctgctgggca ggaagactga ggagggtggg gaccggggcg 1140

ccccagggct ggccctggca cgggtgcggg cggccagcga caccaccaac ggaacaagtt 1200

ccagcaaagg caccagccac agcaaagggc agcggagtgc ctcttcacc taccaccgcc 1260

agcgcaggca tagcgatttc tgtggcccat cccctgcacc cctgcacccc aaacgcagcc 1320

cgacgagcac gggggaggcg gagctgaagg aggagcggct gccaggccgg aaggcgagct 1380

gcagcaccgc ggggagtggg agtcagggc tgccccctc cagcccatg gtcagcagcg 1440

cccacaaccc caacaaggca gagatcccag agcggcgga ggacagcac agcaccacca 1500

acaacctccc tctagcatg atgacctgca gaaacaccta cgtttgcaca gaacgcccgg 1560

gggctgagcg cccgtcactg ttgccaaatg ggaaagaaaa cagctcaggc accccacggg 1620
 tgccccctgc ctccccctcc agtcacagcc tggcaccccc atcagggggag cggagccgcc 1680
 tggcacgagg ttccaccatc cgcagcacct tccatggtgg ccagggtccgg gaccggcggg 1740
 caggggggtgg ggggtgggtgg ggtgtgcaga atggggcccc tgccctctccc aacttgcccc 1800
 atgaggctgc acccctgccc gccggggcggc cccgccccac caccaacctc ttcaccaagc 1860
 tgacctccaa actgacctga agggttaccc tcgatccctc taaacggcag aactctaacc 1920
 gctgtgtttc ggggcgcctc ctgccccagg gatccaagat cagggtcgag acgaacctga 1980
 gagaatcggg ggacctgagg tcacaagttg ccatctacct tgggatcaaa cggaaaccgc 2040
 ccccggtctg ctccgattcc cctggagtgt gaagctgacc agctcgcgcc ctccctgaggc 2100
 cctgatggca gctctgcgcc aggccacagc agccgcccgc tgcgctgcc gccagccaca 2160
 gccgttcctg ctggcctgcc tgcacggggg tgcggggcggg cccgagcccc tgtcccactt 2220
 cgaagtggag gtctgccagc tgccccggcc aggcctgcgg ggagttctct tccgcctgt 2280
 ggcgggcacc gccctggcct tccgcacctc cgtcacccgc atctccaacg acctcgagct 2340
 ctgagccacc acggtcccag ggcccttact ctccctctcc ctgtgcct tcaattctac 2400
 aggaggggaa gggggccagg aggggattct ccttttatca tcacctcagt ttccctgaat 2460
 tataatttgg ggcaaagatt gtcccctctg ctgttctctg gggccgctca gcacagaaga 2520
 aggatgaggg ggctcagcgg ggggagctgg caccttcctg gagcctccag ccagtcctgt 2580
 cctccctcgc cctaccaaga gggcacctga ggagactttg gggacagggc aggggcaggg 2640
 agggaaactg aggaaatctt ccattcctcc caacagctca aaattaggcc ttgggcaggg 2700
 gcaggagag ctgctgagcc taaagactgg agaactctgg ggactgggag tgggggtcag 2760
 agaggcagat tccttccct cccgtccct cagctcaaa ccccaacttc ctgccccagg 2820
 ctggcgcggg gcactttgta caaatccttg taaatacccc acaccctccc ctctgcaaag 2880
 gtctcttgag gagctgccgc tgtcacctac ggtttttaag ttattacacc ccgacctcc 2940
 tcctgtcagc cccctcacct gcagcctgtt gcccaataaa ttaagagag tccccccctc 3000
 cccaatgctg acctaggat tttccttccc tgccctcacc tgcaaagag ttaaagaaga 3060
 ggcgtgggaa tocaggcagt ggtttttcct ttcggagcct cggttttctc atctgcagaa 3120
 tgggagcggg ggggggtggg aggtaaggat ggtcgtggaa gaaggcagga tggaaactcg 3180
 cctcatcccc gagggcccag ttccatatatc gggcccccca ttcattccact cacactccca 3240
 gccaccatgt tacactggac tctaagccac ttcttactcc agtagtaaat ttattcaata 3300
 aacaatcatt gacctatgc tactccatgc caggccaggt gctggacaca gagacatgaa 3360
 gctctgtctg tgggagacag ggattctgac acagacaccg gacaaacct tgtcttgggg 3420

agcccagaag agaaaagtggg caggggtgggg tcattgggga agatgctcta gaggaattaa 3480
 tgctggaatg ggggtgttgaa ggatgagtag gagttagtta ggcattgagt ttgcctctggg 3540
 caaaagccca gaagtgggag tatgtggtat atcttcagag aactgggtaa ttccagtgtg 3600
 gctgctgtgt tgggcatgga tggagaatca gcaagagaaa tgctgtatta ggactaataa 3660
 tccatctacg ctgcttaagc aaaaaggtat ttgttggttt atgttactta atagtccagg 3720
 ggcacctggc ttcaggtagg tttgatccag gcatcaggcc attgcatcta ttttttcagt 3780
 gtaagttgaa ttctagtaat ttttatcaag taagggtctcc tttcctgggtg gcacagatga 3840
 cttcagcagt tagaagtttc tatccctcca gctttctgca gcagaaagac cctcattgtc 3900
 agtttccag caaaagtccc agggcagact ctcatggcc caaatgggcc atgtgatttt 3960
 ctctaaacca atcactgtga ctctagagt gccagactca gagctgcact tagtaggggt 4020
 tcctcaaagg aaggtcaagt gtcatgagca ggagaaaagg catgggagct ggacagatta 4080
 tagtggttga agtctgtgca gtacagaagg gcggagctta ttcacacagc acctttgggg 4140
 ccaaatgaa taagctggac tttctccca tggcactggg gaaccatgga agttcaggga 4200
 acttcaggga agaggcttgg tcaattctg agagcatcct ctgtgctggg gacacagtgg 4260
 taatcaagac agccccaaca ctgccctcat agagctcaca gtccaatgga ggaggcagat 4320
 gtgtcctcag gcagcgactg ggcagggtg gtatagggga gtccagaggt gatgcctgcc 4380
 tcagccaggg agggcttcct ggaggagaag gagccagcta gacatggata ggagtgcgtt 4440
 ttaggcacag caaatggcac atacaagggc caggagcaa gagagaggac aggtcctcaa 4500
 caaatggcat gtgactttgt aagtgtagaa ttgctgtgag gtatggggct agggggtca 4560
 gtagggcctt gaaggttatg gacaggggcc tgggctttct tccaagggca ctgggggagc 4620
 catggcaagg ttgtaggtag ggtagagatg ggcgggtttg tgctatgtgc aggggtggaag 4680
 ggagggaagt tgacaggtca gaagatcagg aaagaggtcg gggctggaca gatggggaga 4740
 gcgcagatag atttaagaga gtccctgtgag gcaaagtggg caggacctgg taacaggtgt 4800
 ctggactgtg gctttggctg gctcagaagg tccccactgg cgtgtgtggt ctatgtagcc 4860
 tctgggtgtg gagctgggat cttcaactgg ggacagtaca gtaaagaaca tcacagc 4917

<210> 23

<211> 3226

<212> DNA

<213> Homo sapiens

<400> 23

gacccggaga agatgtcttc gcggacggtg ctggccccgg gcaacgatcg gaactcggac 60
 acgcatggca ccttgggcag tggcgcctcc tcggacaaag gcccgctctg gtccagccgc 120
 tcaactgggtg cccgttgccg gaactccatc gcctcctgtc ccgaggagca gccccacgtg 180

ggcaactacc gcctgctgag gaccattggg aagggcaact ttgccaaagt caagctggct	240
cggcacatcc tcaactggtcg ggaggttgcc atcaagatta tcgacaaaac ccagctgaat	300
cccagcagcc tgcagaagct gttccgagaa gtccgcacatca tgaagggcct aaaccacccc	360
aacatcgtga agctctttga ggtgattgag actgagaaga cgctgtacct ggtgatggag	420
tacgcaagtg ctggagaagt gtttgactac ctctgtgtcg atggccgcac gaaggagaag	480
gaagctcgag ccaagttccg acagattgtt tccgctgtgc actattgtca ccagaaaaat	540
attgtacaca gggacctgaa ggctgagaac ctcttgctgg atgccgaggg caacatcaag	600
attgctgact ttggcttcag caacgagttc acgctgggat cgaagctgga cacgttctgc	660
gggagcccc catatgccgc cccggagctg tttcagggca agaagtacga cgggcccggag	720
gtggacatct ggagcctggg agtcatcctg tacaccctcg tcagcggctc cctgcccttc	780
gacgggcaca acctcaagga gctgcgggag cgagtactca gagggaagta cggggtccct	840
ttctacatgt caacagactg tgagagcatc ctgcggagat ttttggtgct gaaccagct	900
aaacgctgta ctctcgagca aatcatgaaa gacaaatgga tcaacatcgg ctatgagggt	960
gaggagtga agccatacac agagcccgag gaggacttcg gggacaccaa gagaattgag	1020
gtgatggtgg gtatgggcta cacacgggaa gaaatcaaag agtccttgac cagccagaag	1080
tacaacgaag tgaccgccac ctacctcctg ctgggcagga agactgagga ggggtggggac	1140
egggggcccc cagggctggc cctgggcagg gtgcggggcg ccagcgacac caccaacgga	1200
acaagttcca gcaaaggcac cagccacagc aaagggcagc ggagttcctc ttccacctac	1260
caccgccagc gcaggcatag cgatttctgt ggcaccatccc ctgcacccct gcaccccaaa	1320
cgcagcccga cgagcacggg ggaggcggag ctgaaggagg agcggctgcc aggccggaag	1380
gcgagctgca gcaccgagg gagtgggagt cgagggctgc cccctccag ccccatggtc	1440
agcagcggcc acaaccccaa caaggcagag atcccagagc ggcggaagga cagcacgagc	1500
accccaaca acctccctcc tagcatgatg accgcagaa acacctacgt ttgcacagaa	1560
cgcccggggg ctgagcgccc gtcactgttg ccaaattggga aagaaaacag ctcaggcacc	1620
ccacgggtgc cccctgcctc cccctccagt cacagcctgg ccccccatc aggggagcgg	1680
agccgcctgg cacgcggttc caccatccgc agcaccttc atggtggcca ggtccgggac	1740
cggcgggcag ggggtggggg tgggtggggg gtgcagaatg ggcccctgc ctctcccaca	1800
ctggcccatg aggctgcacc cctgcccgcc gggcggcccc gcccaccac caacctcttc	1860
accaagctga cctccaaact gaccgaagg gtgcagacg aacctgagag aatcggggga	1920
cctgaggtca caagttgcca tctaccttg gatcaaacgg aaaccgcccc ccggtgctc	1980
cgattccctt ggagtgtgaa gctgaccagc tcgcgcctc ctgaggccct gatggcagct	2040

ctgcgccagg ccacagcagc cgcccgtgc cgctgccgcc agccacagcc gttcctgctg 2100
 gcctgcctgc acgggggtgc gggcgggccc gagccctgt occacttcga agtggaggtc 2160
 tgccagctgc cccggccagg cttgcgggga gttctcttcc gccgtgtggc gggcaccgcc 2220
 ctggccttcc gcaccctgt caccgcctc tccaacgacc tcgagctctg agccaccacg 2280
 gtccagggcc cttatcttct cttccttgtc gcttcacttc tacaggaggg gaaggggcca 2340
 gggaggggat tctcccttta tcatcacctc agtttccctg aattatattt gggggcacaag 2400
 attgtccctt ctgctgttct ctggggccgc tcagcacaga agaaggatga gggggctcag 2460
 cggggggagc tggcaccttc ctggagcctc cagccagtcc tgtcctccct cgccctacca 2520
 agagggcacc tgaggagact ttggggacag ggcaggggca gggagggaaa ctgaggaaat 2580
 ctccattcc tcccaacagc tcaaaattag gccttgggca ggggcaggga gagctgctga 2640
 gcctaaagac tggagaatct gggggactgg gagtgggggt cagagaggca gattccttcc 2700
 cctcccgctc cctcacgctc aaacccccac ttcttgcctc aggttgggcg ggggcacttt 2760
 gtacaaatcc ttgtaaatac cccacacctt cccttctgca aaggtctctt gaggagctgc 2820
 cgctgtcacc tacggttttt aagttattac accccgacct tctctctgtc agccccctca 2880
 cgtgcagcct gttgccaat aaatttagga gagtcccccc ctccccaatg ctgaccctag 2940
 gattttcctt ccctgccctc acctgcaaat gagttaaaga agaggcgtgg gaatccaggc 3000
 agtgggtttt cctttcggag cctcgggttt ctcacttgca gaatgggagc ggtgggggtg 3060
 ggaaggtaag gatggtcgtc caagaaggca ggatggaact cggcctcctc cccgaggccc 3120
 cagttcctat atcgggcccc ccattcatcc actcacactc ccagccacca tgttacactg 3180
 gactttaagc catttcttac tccagtagta aatttattca ataaac 3226

<210> 24
 <211> 745
 <212> PRT
 <213> Homo sapiens

<400> 24

Met Ile Arg Gly Arg Asn Ser Ala Thr Ser Ala Asp Glu Gln Pro His
 1 5 10 15

Ile Gly Asn Tyr Arg Leu Leu Lys Thr Ile Gly Lys Gly Asn Phe Ala
 20 25 30

Lys Val Lys Leu Ala Arg His Ile Leu Thr Gly Lys Glu Val Ala Val
 35 40 45

Lys Ile Ile Asp Lys Thr Gln Leu Asn Ser Ser Ser Leu Gln Lys Leu

50		55		60
Phe Arg Glu Val Arg Ile Met Lys Val Leu Asn His Pro Asn Ile Val				
65		70		75
				80
Lys Leu Phe Glu Val Ile Glu Thr Glu Lys Thr Leu Tyr Leu Val Met				
	85		90	
				95
Glu Tyr Ala Ser Gly Gly Glu Val Phe Asp Tyr Leu Val Ala His Gly				
	100		105	110
Arg Met Lys Glu Lys Glu Ala Arg Ala Lys Phe Arg Gln Ile Val Ser				
	115		120	125
Ala Val Gln Tyr Cys His Gln Lys Phe Ile Val His Arg Asp Leu Lys				
	130		135	140
Ala Glu Asn Leu Leu Leu Asp Ala Asp Met Asn Ile Lys Ile Ala Asp				
145		150		155
				160
Phe Gly Phe Ser Asn Glu Phe Thr Phe Gly Asn Lys Leu Asp Thr Phe				
	165		170	
				175
Cys Gly Ser Pro Pro Tyr Ala Ala Pro Glu Leu Phe Gln Gly Lys Lys				
	180		185	190
Tyr Asp Gly Pro Glu Val Asp Val Trp Ser Leu Gly Val Ile Leu Tyr				
	195		200	205
Thr Leu Val Ser Gly Ser Leu Pro Phe Asp Gly Gln Asn Leu Lys Glu				
	210		215	220
Leu Arg Glu Arg Val Leu Arg Gly Lys Tyr Arg Ile Pro Phe Tyr Met				
225		230		235
				240
Ser Thr Asp Cys Glu Asn Leu Leu Lys Lys Phe Leu Ile Leu Asn Pro				
	245		250	255
Ser Lys Arg Gly Thr Leu Glu Gln Ile Met Lys Asp Arg Trp Met Asn				
	260		265	270
Val Gly His Glu Asp Asp Glu Leu Lys Pro Tyr Val Glu Pro Leu Pro				
	275		280	285
Asp Tyr Lys Asp Pro Arg Arg Thr Glu Leu Met Val Ser Met Gly Tyr				
	290		295	300

Thr Arg Glu Glu Ile Gln Asp Ser Leu Val Gly Gln Arg Tyr Asn Glu
 305 310 315 320
 Val Met Ala Thr Tyr Leu Leu Leu Gly Tyr Lys Ser Ser Glu Leu Glu
 325 330 335
 Gly Asp Thr Ile Thr Leu Lys Pro Arg Pro Ser Ala Asp Leu Thr Asn
 340 345 350
 Ser Ser Ala Gln Phe Pro Ser His Lys Val Gln Arg Ser Val Ser Ala
 355 360 365
 Asn Pro Lys Gln Arg Arg Phe Ser Asp Gln Ala Gly Pro Ala Ile Pro
 370 375 380
 Thr Ser Asn Ser Tyr Ser Lys Lys Thr Gln Ser Asn Asn Ala Glu Asn
 385 390 395 400
 Lys Arg Pro Glu Glu Asp Arg Glu Ser Gly Arg Lys Ala Ser Ser Thr
 405 410 415
 Ala Lys Val Pro Ala Ser Pro Leu Pro Gly Leu Glu Arg Lys Lys Thr
 420 425 430
 Thr Pro Thr Pro Ser Thr Asn Ser Val Leu Ser Thr Ser Thr Asn Arg
 435 440 445
 Ser Arg Asn Ser Pro Leu Leu Glu Arg Ala Ser Leu Gly Gln Ala Ser
 450 455 460
 Ile Gln Asn Gly Lys Asp Ser Leu Thr Met Pro Gly Ser Arg Ala Ser
 465 470 475 480
 Thr Ala Ser Ala Ser Ala Ala Val Ser Ala Ala Arg Pro Arg Gln His
 485 490 495
 Gln Lys Ser Met Ser Ala Ser Val His Pro Asn Lys Ala Ser Gly Leu
 500 505 510
 Pro Pro Thr Glu Ser Asn Cys Glu Val Pro Arg Pro Ser Thr Ala Pro
 515 520 525
 Gln Arg Val Pro Val Ala Ser Pro Ser Ala His Asn Ile Ser Ser Ser
 530 535 540
 Gly Gly Ala Pro Asp Arg Thr Asn Phe Pro Arg Gly Val Ser Ser Arg

545 550 555 560
 Ser Thr Phe His Ala Gly Gln Leu Arg Gln Val Arg Asp Gln Gln Asn
 565 570 575
 Leu Pro Tyr Gly Val Thr Pro Ala Ser Pro Ser Gly His Ser Gln Gly
 580 585 590
 Arg Arg Gly Ala Ser Gly Ser Ile Phe Ser Lys Phe Thr Ser Lys Phe
 595 600 605
 Val Arg Arg Asn Leu Asn Glu Pro Glu Ser Lys Asp Arg Val Glu Thr
 610 615 620
 Leu Arg Pro His Val Val Gly Ser Gly Gly Asn Asp Lys Glu Lys Glu
 625 630 635 640
 Glu Phe Arg Glu Ala Lys Pro Arg Ser Leu Arg Phe Thr Trp Ser Met
 645 650 655
 Lys Thr Thr Ser Ser Met Glu Pro Asn Glu Met Met Arg Glu Ile Arg
 660 665 670
 Lys Val Leu Asp Ala Asn Ser Cys Gln Ser Glu Leu His Glu Lys Tyr
 675 680 685
 Met Leu Leu Cys Met His Gly Thr Pro Gly His Glu Asp Phe Val Gln
 690 695 700
 Trp Glu Met Glu Val Cys Lys Leu Pro Arg Leu Ser Leu Asn Gly Val
 705 710 715 720
 Arg Phe Lys Arg Ile Ser Gly Thr Ser Met Ala Phe Lys Asn Ile Ala
 725 730 735
 Ser Lys Ile Ala Asn Glu Leu Lys Leu
 740 745

 <210> 25
 <211> 795
 <212> PRT
 <213> Homo sapiens

 <400> 25
 Met Ser Ala Arg Thr Pro Leu Pro Thr Val Asn Glu Arg Asp Thr Val
 1 5 10 15

Asn His Thr Thr Val Asp Gly Tyr Thr Glu Pro His Ile Gln Pro Thr
 20 25 30

Lys Ser Ser Ser Arg Gln Asn Ile Pro Arg Cys Arg Asn Ser Ile Thr
 35 40 45

Ser Ala Thr Asp Glu Gln Pro His Ile Gly Asn Tyr Arg Leu Gln Lys
 50 55 60

Thr Ile Gly Lys Gly Asn Phe Ala Lys Val Lys Leu Ala Arg His Val
 65 70 75 80

Leu Thr Gly Arg Glu Val Ala Val Lys Ile Ile Asp Lys Thr Gln Leu
 85 90 95

Asn Pro Thr Ser Leu Gln Lys Leu Phe Arg Glu Val Arg Ile Met Lys
 100 105 110

Ile Leu Asn His Pro Asn Ile Val Lys Leu Phe Glu Val Ile Glu Thr
 115 120 125

Glu Lys Thr Leu Tyr Leu Val Met Glu Tyr Ala Ser Gly Gly Glu Val
 130 135 140

Phe Asp Tyr Leu Val Ala His Gly Arg Met Lys Glu Lys Glu Ala Arg
 145 150 155 160

Ala Lys Phe Arg Gln Ile Val Ser Ala Val Gln Tyr Cys His Gln Lys
 165 170 175

Tyr Ile Val His Arg Asp Leu Lys Ala Glu Asn Leu Leu Leu Asp Gly
 180 185 190

Asp Met Asn Ile Lys Ile Ala Asp Phe Gly Phe Ser Asn Glu Phe Thr
 195 200 205

Val Gly Asn Lys Leu Asp Thr Phe Cys Gly Ser Pro Pro Tyr Ala Ala
 210 215 220

Pro Glu Leu Phe Gln Gly Lys Lys Tyr Asp Gly Pro Glu Val Asp Val
 225 230 235 240

Trp Ser Leu Gly Val Ile Leu Tyr Thr Leu Val Ser Gly Ser Leu Pro
 245 250 255

Phe Asp Gly Gln Asn Leu Lys Glu Leu Arg Glu Arg Val Leu Arg Gly
 260 265 270

Lys Tyr Arg Ile Pro Phe Tyr Met Ser Thr Asp Cys Glu Asn Leu Leu
 275 280 285

Lys Lys Leu Leu Val Leu Asn Pro Ile Lys Arg Gly Ser Leu Glu Gln
 290 295 300

Ile Met Lys Asp Arg Trp Met Asn Val Gly His Glu Glu Glu Glu Leu
 305 310 315 320

Lys Pro Tyr Thr Glu Pro Asp Pro Asp Phe Asn Asp Thr Lys Arg Ile
 325 330 335

Asp Ile Met Val Thr Met Gly Phe Ala Arg Asp Glu Ile Asn Asp Ala
 340 345 350

Leu Ile Asn Gln Lys Tyr Asp Glu Val Met Ala Thr Tyr Ile Leu Leu
 355 360 365

Gly Arg Lys Pro Pro Glu Phe Glu Gly Gly Glu Ser Leu Ser Ser Gly
 370 375 380

Asn Leu Cys Gln Arg Ser Arg Pro Ser Ser Asp Leu Asn Asn Ser Thr
 385 390 395 400

Leu Gln Ser Pro Ala His Leu Lys Val Gln Arg Ser Ile Ser Ala Asn
 405 410 415

Gln Lys Gln Arg Arg Phe Ser Asp His Ala Gly Pro Ser Ile Pro Pro
 420 425 430

Ala Val Ser Tyr Thr Lys Arg Pro Gln Ala Asn Ser Val Glu Ser Glu
 435 440 445

Gln Lys Glu Glu Trp Asp Lys Asp Val Ala Arg Lys Leu Gly Ser Thr
 450 455 460

Thr Val Gly Ser Lys Ser Glu Met Thr Ala Ser Pro Leu Val Gly Pro
 465 470 475 480

Glu Arg Lys Lys Ser Ser Thr Ile Pro Ser Asn Asn Val Tyr Ser Gly
 485 490 495

Gly Ser Met Ala Arg Arg Asn Thr Tyr Val Cys Glu Arg Thr Thr Asp
 500 505 510

Arg Tyr Val Ala Leu Gln Asn Gly Lys Asp Ser Ser Leu Thr Glu Met
515 520 525

Ser Val Ser Ser Ile Ser Ser Ala Gly Ser Ser Val Ala Ser Ala Val
530 535 540

Pro Ser Ala Arg Pro Arg His Gln Lys Ser Met Ser Thr Ser Gly His
545 550 555 560

Pro Ile Lys Val Thr Leu Pro Thr Ile Lys Asp Gly Ser Glu Ala Tyr
565 570 575

Arg Pro Gly Thr Thr Gln Arg Val Pro Ala Ala Ser Pro Ser Ala His
580 585 590

Ser Ile Ser Thr Ala Thr Pro Asp Arg Thr Arg Phe Pro Arg Gly Ser
595 600 605

Ser Ser Arg Ser Thr Phe His Gly Glu Gln Leu Arg Glu Arg Arg Ser
610 615 620

Val Ala Tyr Asn Gly Pro Pro Ala Ser Pro Ser His Glu Thr Gly Ala
625 630 635 640

Phe Ala His Ala Arg Arg Gly Thr Ser Thr Gly Ile Ile Ser Lys Ile
645 650 655

Thr Ser Lys Phe Val Arg Arg Asp Pro Ser Glu Gly Glu Ala Ser Gly
660 665 670

Arg Thr Asp Thr Ser Arg Ser Thr Ser Gly Glu Pro Lys Glu Arg Asp
675 680 685

Lys Glu Glu Gly Lys Asp Ser Lys Pro Arg Ser Leu Arg Phe Thr Trp
690 695 700

Ser Met Lys Thr Thr Ser Ser Met Asp Pro Asn Asp Met Met Arg Glu
705 710 715 720

Ile Arg Lys Val Leu Asp Ala Asn Asn Cys Asp Tyr Glu Gln Lys Glu
725 730 735

Arg Phe Leu Leu Phe Cys Val His Gly Asp Ala Arg Gln Asp Ser Leu
740 745 750

Val Gln Trp Glu Met Glu Val Cys Lys Leu Pro Arg Leu Ser Leu Asn
755 760 765

Gly Val Arg Phe Lys Arg Ile Ser Gly Thr Ser Ile Ala Phe Lys Asn
 770 775 780

Ile Ala Ser Lys Ile Ala Asn Glu Leu Lys Leu
 785 790 795

<210> 26
 <211> 729
 <212> PRT
 <213> Homo sapiens

<400> 26

Met Ser Thr Arg Thr Pro Leu Pro Thr Val Asn Glu Arg Asp Thr Glu
 1 5 10 15

Asn His Thr Ser His Gly Asp Gly Arg Gln Glu Val Thr Ser Arg Thr
 20 25 30

Ser Arg Ser Gly Ala Arg Cys Arg Asn Ser Ile Ala Ser Cys Ala Asp
 35 40 45

Glu Gln Pro His Ile Gly Asn Tyr Arg Leu Leu Lys Thr Ile Gly Lys
 50 55 60

Gly Asn Phe Ala Lys Val Lys Leu Ala Arg His Ile Leu Thr Gly Arg
 65 70 75 80

Glu Val Ala Ile Lys Ile Ile Asp Lys Thr Gln Leu Asn Pro Thr Ser
 85 90 95

Leu Gln Lys Leu Phe Arg Glu Val Arg Ile Met Lys Ile Leu Asn His
 100 105 110

Pro Asn Ile Val Lys Leu Phe Glu Val Ile Glu Thr Glu Lys Thr Leu
 115 120 125

Tyr Leu Ile Met Glu Tyr Ala Ser Gly Gly Glu Val Phe Asp Tyr Leu
 130 135 140

Val Ala His Gly Arg Met Lys Glu Lys Glu Ala Arg Ser Lys Phe Arg
 145 150 155 160

Gln Ile Val Ser Ala Val Gln Tyr Cys His Gln Lys Arg Ile Val His
 165 170 175

Arg Asp Leu Lys Ala Glu Asn Leu Leu Leu Asp Ala Asp Met Asn Ile

180	185	190
Lys Ile Ala Asp Phe Gly Phe Ser Asn Glu Phe Thr Val Gly Gly Lys 195 200 205		
Leu Asp Thr Phe Cys Gly Ser Pro Pro Tyr Ala Ala Pro Glu Leu Phe 210 215 220		
Gln Gly Lys Lys Tyr Asp Gly Pro Glu Val Asp Val Trp Ser Leu Gly 225 230 235 240		
Val Ile Leu Tyr Thr Leu Val Ser Gly Ser Leu Pro Phe Asp Gly Gln 245 250 255		
Asn Leu Lys Glu Leu Arg Glu Arg Val Leu Arg Gly Lys Tyr Arg Ile 260 265 270		
Pro Phe Tyr Met Ser Thr Asp Cys Glu Asn Leu Leu Lys Arg Phe Leu 275 280 285		
Val Leu Asn Pro Ile Lys Arg Gly Thr Leu Glu Gln Ile Met Lys Asp 290 295 300		
Arg Trp Ile Asn Ala Gly His Glu Glu Asp Glu Leu Lys Pro Phe Val 305 310 315 320		
Glu Pro Glu Leu Asp Ile Ser Asp Gln Lys Arg Ile Asp Ile Met Val 325 330 335		
Gly Met Gly Tyr Ser Gln Glu Glu Ile Gln Glu Ser Leu Ser Lys Met 340 345 350		
Lys Tyr Asp Glu Ile Thr Ala Thr Tyr Leu Leu Leu Gly Arg Lys Ser 355 360 365		
Ser Glu Leu Asp Ala Ser Asp Ser Ser Ser Ser Ser Asn Leu Ser Leu 370 375 380		
Ala Lys Val Arg Pro Ser Ser Asp Leu Asn Asn Ser Thr Gly Gln Ser 385 390 395 400		
Pro His His Lys Val Gln Arg Ser Val Ser Ser Ser Gln Lys Gln Arg 405 410 415		
Arg Tyr Ser Asp His Ala Gly Pro Ala Ile Pro Ser Val Val Ala Tyr 420 425 430		

Pro Lys Arg Ser Gln Thr Ser Thr Ala Asp Gly Asp Leu Lys Glu Asp
 435 440 445

Gly Ile Ser Ser Arg Lys Ser Ser Gly Ser Ala Val Gly Gly Lys Gly
 450 455 460

Ile Ala Pro Ala Ser Pro Met Leu Gly Asn Ala Ser Asn Pro Asn Lys
 465 470 475 480

Ala Asp Ile Pro Glu Arg Lys Lys Ser Ser Thr Val Pro Ser Ser Asn
 485 490 495

Thr Ala Ser Gly Gly Met Thr Arg Arg Asn Thr Tyr Val Cys Ser Glu
 500 505 510

Arg Thr Thr Ala Asp Arg His Ser Val Ile Gln Asn Gly Lys Glu Asn
 515 520 525

Ser Thr Ile Pro Asp Gln Arg Thr Pro Val Ala Ser Thr His Ser Ile
 530 535 540

Ser Ser Ala Ala Thr Pro Asp Arg Ile Arg Phe Pro Arg Gly Thr Ala
 545 550 555 560

Ser Arg Ser Thr Phe His Gly Gln Pro Arg Glu Arg Arg Thr Ala Thr
 565 570 575

Tyr Asn Gly Pro Pro Ala Ser Pro Ser Leu Ser His Glu Ala Thr Pro
 580 585 590

Leu Ser Gln Thr Arg Ser Arg Gly Ser Thr Asn Leu Phe Ser Lys Leu
 595 600 605

Thr Ser Lys Leu Thr Arg Ser Arg Asn Val Ser Ala Glu Gln Lys Asp
 610 615 620

Glu Asn Lys Glu Ala Lys Pro Arg Ser Leu Arg Phe Thr Trp Ser Met
 625 630 635 640

Lys Thr Thr Ser Ser Met Asp Pro Gly Asp Met Met Arg Glu Ile Arg
 645 650 655

Lys Val Leu Asp Ala Asn Asn Cys Asp Tyr Glu Gln Arg Glu Arg Phe
 660 665 670

Leu Leu Phe Cys Val His Gly Asp Gly His Ala Glu Asn Leu Val Gln

675

680

685

Trp Glu Met Glu Val Cys Lys Leu Pro Arg Leu Ser Leu Asn Gly Val
690 695 700

Arg Phe Lys Arg Ile Ser Gly Thr Ser Ile Ala Phe Lys Asn Ile Ala
705 710 715 720

Ser Lys Ile Ala Asn Glu Leu Lys Leu
725

<210> 27
<211> 713
<212> PRT
<213> Homo sapiens

<400> 27

Met Ser Thr Arg Thr Pro Leu Pro Thr Val Asn Glu Arg Asp Thr Glu
1 5 10 15

Asn His Thr Ser His Gly Asp Gly Arg Gln Glu Val Thr Ser Arg Thr
20 25 30

Ser Arg Ser Gly Ala Arg Cys Arg Asn Ser Ile Ala Ser Cys Ala Asp
35 40 45

Glu Gln Pro His Ile Gly Asn Tyr Arg Leu Leu Lys Thr Ile Gly Lys
50 55 60

Gly Asn Phe Ala Lys Val Lys Leu Ala Arg His Ile Leu Thr Gly Arg
65 70 75 80

Glu Val Ala Ile Lys Ile Ile Asp Lys Thr Gln Leu Asn Pro Thr Ser
85 90 95

Leu Gln Lys Leu Phe Arg Glu Val Arg Ile Met Lys Ile Leu Asn His
100 105 110

Pro Asn Ile Val Lys Leu Phe Glu Val Ile Glu Thr Gln Lys Thr Leu
115 120 125

Tyr Leu Ile Met Glu Tyr Ala Ser Gly Gly Lys Val Phe Asp Tyr Leu
130 135 140

Val Ala His Gly Arg Met Lys Glu Lys Glu Ala Arg Ser Lys Phe Arg
145 150 155 160

Gln Ile Val Ser Ala Val Gln Tyr Cys His Gln Lys Arg Ile Val His
 165 170 175
 Arg Asp Leu Lys Ala Glu Asn Leu Leu Asp Ala Asp Met Asn Ile
 180 185 190
 Lys Ile Ala Asp Phe Gly Phe Ser Asn Glu Phe Thr Val Gly Gly Lys
 195 200 205
 Leu Asp Thr Phe Cys Gly Ser Pro Pro Tyr Ala Ala Pro Glu Leu Phe
 210 215 220
 Gln Gly Lys Lys Tyr Asp Gly Pro Glu Val Asp Val Trp Ser Leu Gly
 225 230 235 240
 Val Ile Leu Tyr Thr Leu Val Ser Gly Ser Leu Pro Phe Asp Gly Gln
 245 250 255
 Asn Leu Lys Glu Leu Arg Glu Arg Val Leu Arg Gly Lys Tyr Arg Ile
 260 265 270
 Pro Phe Tyr Met Ser Thr Asp Cys Glu Asn Leu Leu Lys Arg Phe Leu
 275 280 285
 Val Leu Asn Pro Ile Lys Arg Gly Thr Leu Glu Gln Ile Met Lys Asp
 290 295 300
 Arg Trp Ile Asn Ala Gly His Glu Glu Asp Glu Leu Lys Pro Phe Val
 305 310 315 320
 Glu Pro Glu Leu Asp Ile Ser Asp Gln Lys Arg Ile Asp Ile Met Val
 325 330 335
 Gly Met Gly Tyr Ser Gln Glu Glu Ile Gln Glu Ser Leu Ser Lys Met
 340 345 350
 Lys Tyr Asp Glu Ile Thr Ala Thr Tyr Leu Leu Leu Gly Arg Lys Ser
 355 360 365
 Ser Glu Val Arg Pro Ser Ser Asp Leu Asn Asn Ser Thr Gly Gln Ser
 370 375 380
 Pro His His Lys Val Gln Arg Ser Val Ser Ser Ser Gln Lys Gln Arg
 385 390 395 400
 Arg Tyr Ser Asp His Ala Gly Pro Gly Ile Pro Ser Val Val Ala Tyr
 405 410 415